Yale University, New Haven, Connecticut

2004 EPA STAR Graduate Fellowship Conference Next Generation Scientists—Next Opportunities

# Investigating the Ecological Impacts of Anadromous Alewife Restoration in New England

## **Environmental Issue**

•Dam removal and fish ladder construction projects are underway on many coastal New England rivers and streams

•These projects aim to restore anadromous (migratory) fish species, including the historically abundant alewife

•Restored access to historical spawning habitat will have unexplored ecological consequences

•One potential hurdle to anadromous alewife restoration is the presence of landlocked alewife populations in many lakes



### Anadromous Alewife Life Cycle

•Spring migrations bring adults from the Atlantic Ocean into coastal rivers and lakes where spawning takes place

•Larvae feed on zooplankton in freshwater and grow throughout the summer

•In early fall, the juveniles migrate to the ocean where they grow to sexual maturity in 3-4 years

 Mature adults return to their natal streams to spawn

 Spawners import marine nutrients into freshwater systems and juveniles may shape zooplankton assemblages through intense predation

### Abstract

Populations of anadromous and landlocked alewives (Alosa pseudoharengus) will be studied in order to understand the evolutionary origin and ecological consequences of life history variation in this species. From a management perspective, this study aims to understand the implications of anadromous alewife restoration at both the population and community levels.

The Ecological Consequences of Life History Variation Alewives express anadromous and landlocked life histories. Whereas anadromous alewives migrate between fresh and salt water. landlocked alewives live their entire lives in freshwater lakes and are important pelagic (open water) predators of zooplankton. Because of differences in predation pressure and resource availability. landlocked alewives are smaller and younger at maturity than anadromous alewives.



Freshwater zooplankton are prey for landlocked alewives and for iuvenile anadromous alewives. Alewife predation can shift zooplankton dominance from large bodied species such as Daphnia (top) to small bodied species such as Bosmina (bottom).

### **Kev Questions:**

•Do anadromous and landlocked alewives differ in their ecological roles?

What changes will anadromous alewife recovery bring about in lakes with/without landlocked alewives? Will landlocked alewife populations be a barrier to anadromous alewife recovery?

# **Scientific Approach**



 Anadromous and landlocked populations are sampled to estimate population densities, age-specific survival and reproductive rates, and individual growth rates. These population parameters are input into a model that will be used to predict population growth or decline under different recovery scenarios.

1. Growth, Survival, Reproduction

2. Genetic and Morphological Variation

•Genetic and morphological differences between populations are estimated. These differences may underlie any ecological differences between the two life histories and are useful for inferring the evolutionary origin of landlocked populations.



3. Experimental Manipulations

•The presence of anadromous and landlocked juveniles are manipulated in lake mesocosm experiments which examine the ecological effects of anadromous alewife restoration. Alewife growth and survival, zooplankton diversity, and algae abundance are measured over the duration of the experiment.





