

Cardiovascular Disease

The broad category of cardiovascular disease (CVD) includes any disease involving the heart and blood vessels. Coronary heart disease, cerebrovascular disease (commonly known as stroke), and hypertension are among the major cardiovascular diseases (American Heart Association, 2020). In addition to being a major risk factor for heart disease and stroke, hypertension is a commonly diagnosed disease that can also lead to kidney damage and other health problems. Obesity, physical inactivity, and sodium intake are all important risk factors for hypertension (American Heart Association, 2017). Since 1900, CVD has been the leading cause of death in the U.S. every year except 1918 (American Heart Association, 2016) ([General Mortality](#) indicator). The U.S. age-adjusted death rate for CVD reached a peak in 1950. Between 1950 and 1999, the age-adjusted death rate for CVD declined 60 percent (CDC, 1999). The major risk factors for CVD include tobacco use, high blood pressure, high blood cholesterol, diabetes, physical inactivity, poor nutrition, and being overweight or obese (American Heart Association, 2020).

Environmental exposures may also play a role in CVD morbidity and mortality independent of other risk factors. However, susceptible populations such as the elderly and other high-risk populations may be most impacted. For example, studies have shown exposure to ambient airborne particulate matter could be associated with increased hospitalizations and mortality among older individuals, largely due to cardiopulmonary and cardiovascular disease (American Lung Association, 2020; Brook et al., 2010; U.S. EPA, 2019, 2020). Secondhand smoke (SHS) may also contribute to CVD. Although the smoke to which a nonsmoker is exposed is less concentrated than that inhaled by smokers, research has demonstrated increased cardiovascular-related health risks associated with SHS (HHS, 2014; State of California, 2005).

This indicator presents U.S. adult (age 18 and older) prevalence rates for heart disease (all types), coronary heart disease, stroke, and hypertension; and death rates for CVD as a whole as well as specifically for coronary heart disease (including myocardial infarction), stroke, and hypertension. CVD prevalence data were compiled between 2002 and 2018 from the National Health Interview Survey (NHIS), conducted by the Centers for Disease Control and Prevention's (CDC's) National Center for Health Statistics (NCHS). The NHIS is the principal source of information on the health of the civilian non-institutionalized population of the U.S. and since 1960 has been one of the major data collection programs of NCHS. CVD prevalence is based on the number of adults who reported that they had ever been told by a doctor or other health practitioner that they had a specified CVD.

Mortality data were compiled using the National Vital Statistics System (NVSS), maintained by NCHS. Total mortality data were compiled between 1979 and 2018. Because of the change of reporting requirements for race and ethnicity, data by demographic group (i.e., age, ethnicity, race, and sex) were compiled for the years 1999 to 2018, when data are available to compare more races and ethnicities. The NVSS registers virtually all deaths and births nationwide, with data coverage from 1933 to 2018. The NVSS collects data from 57 vital registration jurisdictions, including all 50 states, the District of Columbia, New York City, and five territories (NCHS, 2020). Data were queried and compared separately for years 1979-1998 and those 1999 onward because the NVSS uses different codes to specify causes of death for these two time periods: the International Classification of Diseases 9th Revision (ICD-9) codes for 1979-1998 and the International Classification of Diseases 10th Revision (ICD-10) codes beginning in 1999.

What the Data Show

CVD Prevalence

Among adults 18 years and older, the age-adjusted prevalence of coronary heart disease and stroke between 2002 and 2018 has remained essentially the same. In contrast, the prevalence of hypertension showed a general increase from 2002 to 2010, but has remained about the same since then (Exhibit 1).

Sex, race, ethnicity, and age differences in CVD prevalence exist. The age-adjusted prevalence of coronary heart disease is consistently higher among males than among females (74 cases per 1,000 for men compared with 41 cases per 1,000 for women in 2018). The prevalence of hypertension and stroke varies by year, with rates among women higher some years while other years' rates were higher among men. As would be expected, the prevalence of heart disease (all types), coronary heart disease, hypertension, and stroke increase as people age, with those aged 18 to 44 years having the lowest prevalence and those 75 years and older having the highest prevalence of CVD (Exhibit 1).

In 2018, the prevalence of coronary heart disease across races was highest among American Indians and Alaska Natives (86 cases per 1,000), followed by Whites (57 cases per 1,000), Blacks or African Americans (54 cases per 1,000), and Asians (44 cases per 1,000). Between 2002 and 2018, Blacks or African Americans consistently had the highest prevalence of hypertension (322 cases per 1,000 in 2018). Asians consistently had the lowest prevalence of hypertension (219 cases per 1,000 in 2018) among the racial groups reported. Asians also had the lowest prevalence of stroke except for four years (2002, 2007, 2011, and 2018) when Whites had the lowest rates (26 cases per 1,000 in 2018). When taking only ethnicity into account, the Hispanic or Latino population had a consistently lower prevalence of heart disease (all types), coronary heart disease, and hypertension from 2002-2018 (except for coronary heart disease in 2016), but rates fluctuated for stroke, when compared with the non-Hispanic or Latino population. In 2018, prevalence in Hispanics or Latinos was lower than in non-Hispanics or Latinos for coronary heart disease (51 versus 57 cases per 1,000, respectively), hypertension (237 versus 251 cases per 1,000, respectively), and stroke (25 cases versus 29 cases per 1,000) (Exhibit 1).

CVD Mortality

In 1979, the national age-adjusted CVD death rate (all types) was 531.0 per 100,000 compared to a rate of 352.0 per 100,000 in 1998 (Exhibit 2). This decline continued from 1999 to 2018, with the rate dropping from 349.3 per 100,000 in 1999 to 215.8 per 100,000 in 2018. Death rates for coronary heart disease, stroke, and myocardial infarction—subcategories of CVD—all declined between 1979 and 1998. The age-adjusted coronary heart disease death rate ranged from 339.2 per 100,000 in 1979 to 197.1 per 100,000 in 1998. The age-adjusted death rate for myocardial infarction ranged from 157.9 in 1979 to 76.0 per 100,000 in 1998. The decline in mortality from coronary heart disease and myocardial infarction continued to be observed between 1999 and 2018. For stroke mortality, the age-adjusted rate ranged from 97.1 per 100,000 in 1979 to 59.3 per 100,000 in 1998. A decline in stroke mortality was also seen from 1999 through 2013, but then slightly increased in 2014 and 2015 and remained relatively steady from 2016 onward. The age-adjusted death rates for coronary heart disease, stroke, and myocardial infarction in 2018 were 90.9, 37.1, and 27.0 per 100,000, respectively, compared to 194.6, 61.6, and 73.2 per 100,000, respectively, in 1999. In contrast, mortality attributed to hypertension fluctuated from 1979 to 1998, with a slight decrease overall from 17.5 per 100,000 in 1979 to 16.6 per 100,000 in 1998. From 1999 to 2018, mortality from hypertension increased overall from 15.8 per 100,000 in 1999 to 24.0 per 100,000 in 2018 (Exhibit 2).

Differences exist in CVD death rates among age, sex, ethnicity, and race groups. For example, in 2018, those aged 65 and older had the highest CVD (all types), coronary heart disease, stroke, hypertension, and myocardial infarction mortality (1,383.8, 573.8, 251.7, 143.5, and 161.8 per 100,000, respectively). Also in 2018, the age-adjusted CVD (all types), coronary heart disease, stroke, hypertension, and myocardial infarction death rates for those 45 to 64 years of age were 152.1, 73.1, 19.4, 21.3, and 26.0 per 100,000, respectively. Notable differences in CVD (all types), coronary heart disease, and myocardial infarction death rates exist between males and females, but not for stroke mortality or hypertension. CVD (all types), coronary heart disease, and myocardial infarction mortality among males in 2018 was 261.9, 124.5, and 36.4 per 100,000, respectively, compared to 177.5, 64.0, and 19.3 per 100,000, respectively, for females. From 1999 to 2018, Blacks or African Americans and non-Hispanics or Latinos had the highest death rates for the five CVD-related diseases among all reported races and ethnicities, respectively. Also, from 1999 to 2018, Asians or Pacific Islanders had the lowest death rates for CVD (all types), coronary heart disease, and myocardial infarction; American Indians and Alaska Natives had the lowest death rates throughout this time period for stroke. Asians or Pacific Islanders had the lowest death rates for hypertension in 2005, 2006, 2010, and 2012-2018, while American Indians and Alaska Natives had the lowest death rates for the other years during the 1999-2018 period (Exhibit 3).

Both coronary heart disease and stroke mortality have been declining over time in each of the 10 EPA Regions (Exhibits 4 and 5, respectively). In 1979, coronary heart disease and stroke age-adjusted death rates ranged from 285.6 (Region 10) to 401.9 (Region 2) per 100,000 and 80.3 (Region 2) to 111.4 (Region 4) per 100,000, respectively. In 1998, coronary heart disease and stroke death rates ranged from 145.6 (Region 8) to 233.2 (Region 2) per 100,000 and 43.2 (Region 2) to 68.5 (Region 10) per 100,000, respectively. The overall decreases in coronary heart disease and stroke mortality also continue in the 1999-2018 period across all

EPA regions. In 1999, coronary heart disease and stroke age-adjusted death rates ranged from 140.4 (Region 8) to 234.8 (Region 2) per 100,000 and 43.8 (Region 2) to 72.8 (Region 10) per 100,000, respectively. In 2018, coronary heart disease and stroke death rates ranged from 71.6 (Region 8) to 108.4 (Region 2) per 100,000 and 26.0 (Region 2) to 42.7 (Region 4) per 100,000, respectively.

Limitations

- Prevalence data reported in the NHIS are based on self-reported responses to specific questions pertaining to CVD-related illnesses, and are subject to the biases associated with self-reported data. Self-reported data can underestimate the disease prevalence being measured if, for whatever reason, the respondent is not fully aware of his/her condition.
- CVD death rates are based on underlying cause of death as entered on a death certificate by a physician, medical examiner, or coroner. Some individuals may have had competing causes of death. When more than one cause or condition is entered by the physician, medical examiner, or coroner, the underlying cause is determined by the sequence of conditions on the certificate, provisions of the ICD [International Classification of Diseases], and associated selection rules and modifications (CDC, 2020b). Consequently, some misclassification of reported mortality might occur in individuals with competing causes of death, as well as the possible underreporting of CVD as the cause of death.
- The International Classification of Diseases 9th Revision (ICD-9) codes were used to specify underlying cause of death for years 1979-1998. Beginning in 1999, cause of death is specified with the International Classification of Diseases 10th Revision (ICD-10) codes. The two revisions differ substantially, and to prevent confusion about the significance of any specific disease code, data queries are separate.

Data Sources

CVD prevalence data were obtained from annual reports and standalone tables published by NCHS (NCHS, 2004, 2005, 2006a,b, 2007, 2009a,b, 2010, 2012a,b, 2014, 2015a,b, 2016, 2018a,b, 2019), which summarize health statistics compiled from the NHIS (<https://www.cdc.gov/nchs/products/series.htm>). CVD mortality statistics for 1999 to 2018 were obtained from CDC's Underlying Cause of Death database, accessed through CDC's Wide-ranging Online Data for Epidemiologic Research (WONDER) (CDC, 2020a) (<https://wonder.cdc.gov/ucd-icd10.html>). Mortality statistics for 1979-1998 in Exhibits 2, 4, and 5 came from the CDC WONDER Compressed Mortality Database (CDC, 2011) (<https://wonder.cdc.gov/mortSQL.html>). EPA Regional mortality statistics were generated by combining and age-adjusting state-by-state totals for each EPA Region using data from CDC WONDER.

References

- American Heart Association. 2020. Heart disease and stroke statistics—2020 update: A report from the American Heart Association. *Circulation Journal of the American Heart Association* (141):e139-e596. <https://www.ahajournals.org/doi/10.1161/CIR.0000000000000757>.
- American Heart Association. 2017. Know your risk factors for high blood pressure. Last reviewed December 2017. Accessed July 9, 2020. <https://www.heart.org/en/health-topics/high-blood-pressure/why-high-blood-pressure-is-a-silent-killer/know-your-risk-factors-for-high-blood-pressure>.
- American Heart Association. 2016. Heart disease and stroke statistics—2016 update: A report from the American Heart Association. *Circulation Journal of the American Heart Association* (133):e38-e360. <https://www.ahajournals.org/doi/epub/10.1161/CIR.0000000000000350>.
- American Lung Association. 2020. State of the air 2020. <https://www.stateoftheair.org/assets/SOTA-2020.pdf> (PDF) (162 pp, 1.9MB).
- Brook, R.D., S. Rajagopalan, C.A. Pope, III, J.R. Brook, A. Bhatnagar, A.V. Diez-Roux, et al. 2010. Particulate matter air pollution and cardiovascular disease: An update to the scientific statement from the American Heart Association. *Circulation* 121:2331-2378. <https://ahajournals.org/doi/full/10.1161/cir.0b013e3181dbee1>.
- CDC (Centers for Disease Control and Prevention). 2020a. Wide-ranging OnLine Data for Epidemiologic Research (WONDER): Underlying cause of death, 1999-2018. Accessed July 2020. <https://wonder.cdc.gov/ucd-icd10.html>.
- CDC. 2020b. Wide-ranging OnLine Data for Epidemiologic Research (WONDER): Underlying cause of death, 1999-2018—help page. Accessed August 6, 2020. <https://wonder.cdc.gov/wonder/help/ucd.html>.
- CDC. 2011. CDC Wide-ranging OnLine Data for Epidemiologic Research (WONDER). Compressed mortality file, underlying cause of death. 1979-1998 (with ICD 9 codes). Accessed April 2012. <https://wonder.cdc.gov/mortSQL.html>.
- CDC. 1999. Decline in deaths from heart disease and stroke, United States, 1990-1999. Washington, DC.
- HHS (U.S. Department of Health and Human Services). 2014. The health consequences of smoking—50 years of progress: A report of the surgeon general. Atlanta, GA: U.S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health. https://www.ncbi.nlm.nih.gov/books/NBK179276/pdf/Bookshelf_NBK179276.pdf (PDF) (1081 pp, 36MB).
- NCHS (National Center for Health Statistics). 2020. National Vital Statistics System fact sheet. April 2020. https://www.cdc.gov/nchs/data/factsheets/factsheet_NVSS.pdf (PDF) (2 pp, 193K).
- NCHS. 2019. Tables of summary health statistics for U.S. adults: 2018 National Health Interview Survey. Table A-1. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2018_SHS_Table_A-1.pdf (PDF) (9 pp, 137K).
- NCHS. 2018a. Tables of summary health statistics for U.S. adults: 2017 National Health Interview Survey. Table A-1. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2017_SHS_Table_A-1.pdf (PDF) (9 pp, 143K).
- NCHS. 2018b. Tables of summary health statistics for U.S. adults: 2016 National Health Interview Survey. Table A-1. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2016_SHS_Table_A-1.pdf (PDF) (9 pp, 143K).
- NCHS. 2016. Tables of summary health statistics for U.S. adults: 2015 National Health Interview Survey. Table A-1. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2015_SHS_Table_A-1.pdf (PDF) (9 pp, 189K).
- NCHS. 2015a. Tables of summary health statistics for U.S. adults: 2014 National Health Interview Survey. Table A-1. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2014_SHS_Table_A-1.pdf (PDF) (9 pp, 178K).
- NCHS. 2015b. Tables of summary health statistics for U.S. adults: 2013 National Health Interview Survey. Table A-1. https://ftp.cdc.gov/pub/Health_Statistics/NCHS/NHIS/SHS/2013_SHS_Table_A-1.pdf (PDF) (9 pp, 119K).
- NCHS. 2014. Summary health statistics for U.S. adults: National Health Interview Survey, 2012. *Vital Health Stat.* 10(260). https://www.cdc.gov/nchs/data/series/sr_10/sr10_260.pdf (PDF) (171 pp, 3.5MB).
- NCHS. 2012a. Summary health statistics for U.S. adults: National Health Interview Survey, 2011. *Vital Health Stat.* 10(256). https://www.cdc.gov/nchs/data/series/sr_10/sr10_256.pdf (PDF) (218 pp, 3.1MB).
- NCHS. 2012b. Summary health statistics for U.S. adults: National Health Interview Survey, 2010. *Vital Health Stat.* 10(252). https://www.cdc.gov/nchs/data/series/sr_10/sr10_252.pdf (PDF) (217 pp, 3.1MB).
- NCHS. 2010. Summary health statistics for U.S. adults: National Health Interview Survey, 2009. *Vital Health Stat.* 10(249). https://www.cdc.gov/nchs/data/series/sr_10/sr10_249.pdf (PDF) (217 pp, 354MB).
- NCHS. 2009a. Summary health statistics for U.S. adults: National Health Interview Survey, 2008. *Vital Health Stat.* 10(242).

https://www.cdc.gov/nchs/data/series/sr_10/sr10_242.pdf (PDF) (167 pp, 2.6MB).

NCHS. 2009b. Summary health statistics for U.S. adults: National Health Interview Survey, 2007. Vital Health Stat. 10(240). https://www.cdc.gov/nchs/data/series/sr_10/sr10_240.pdf (PDF) (169 pp, 8.4MB).

NCHS. 2007. Summary health statistics for U.S. adults: National Health Interview Survey, 2006. Vital Health Stat. 10(235). https://www.cdc.gov/nchs/data/series/sr_10/sr10_235.pdf (PDF) (163 pp, 3.6MB).

NCHS. 2006a. Summary health statistics for U.S. adults: National Health Interview Survey, 2005. Vital Health Stat. 10(232). https://www.cdc.gov/nchs/data/series/sr_10/sr10_232.pdf (PDF) (163 pp, 3.7MB).

NCHS. 2006b. Summary health statistics for U.S. adults: National Health Interview Survey, 2004. Vital Health Stat. 10(228). https://www.cdc.gov/nchs/data/series/sr_10/sr10_228.pdf (PDF) (164 pp, 3.8MB).

NCHS. 2005. Summary health statistics for U.S. adults: National Health Interview Survey, 2003. Vital Health Stat. 10(225). https://www.cdc.gov/nchs/data/series/sr_10/sr10_225.pdf (PDF) (161 pp, 3.8MB).

NCHS. 2004. Summary health statistics for U.S. adults: National Health Interview Survey, 2002. Vital Health Stat. 10(222). https://www.cdc.gov/nchs/data/series/sr_10/sr10_222.pdf (PDF) (161 pp, 4.0MB).


State of California. 2005. Proposed identification of environmental tobacco smoke as a toxic air contaminant. Part B: Health effects assessment for environmental tobacco smoke. As approved by the Scientific Review Panel on June 24, 2005. California Environmental Protection Agency, Office of Environmental Health Hazard Assessment. <https://www.arb.ca.gov/regact/ets2006/app3partb.pdf> (PDF) (528 pp, 4.0MB).

U.S. EPA (United States Environmental Protection Agency). 2020. Integrated science assessment for ozone and related photochemical oxidants. EPA 600/R-20/012. Research Triangle Park, NC. <https://cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=348522>.

U.S. EPA. 2019. Integrated science assessment for particulate matter. EPA/600/R-19/188. Research Triangle Park, NC. <https://cfpub.epa.gov/ncea/isa/recordisplay.cfm?deid=347534>.

Exhibit 1. Age-adjusted cardiovascular disease prevalence in U.S. adults (age 18 and older) by sex, race, ethnicity, and age group, 2002–2018

All cases




Rates are age-adjusted to the 2000 U.S. standard population.

Information on the statistical significance of the trends in this exhibit is not presented here. For more information about uncertainty, variability, and statistical analysis, view the technical documentation for this indicator.

Data source: NCHS, 2004, 2005, 2006a,b, 2007, 2009a,b, 2010, 2012a,b, 2014, 2015a,b, 2016, 2018a,b, 2019

Visit <https://www.epa.gov/roe> to see the full exhibit.

Exhibit 2. Age-adjusted cardiovascular disease death rates in the U.S., 1979-2018



Due to differences in the ICD system used for classifying mortalities, data from 1979-1998 should not be directly compared with data from 1999-2018 [ICD-9 codes: 390-434, 436-448 (1979-1998); ICD-10 codes: I00-I78 (1999-2018)].


Rates are age-adjusted to the 2000 U.S. standard population.

Information on the statistical significance of the trends in this exhibit is not presented here. For more information about uncertainty, variability, and statistical analysis, view the technical documentation for this indicator.

Data source: CDC, 2011, 2020a

Exhibit 3. Age-adjusted cardiovascular disease death rates in the U.S. by sex, race, ethnicity, and age group, 1999-2018

Heart disease (all types)



Rates are age-adjusted to the 2000 U.S. standard population.


Age-adjusted death rates for American Indian/Alaska Natives, Asians or Pacific Islanders, and Hispanics or Latinos are underestimates resulting from an underreporting on death certificates and undercounts in censuses for these groups.

Information on the statistical significance of the trends in this exhibit is not presented here. For more information about uncertainty, variability, and statistical analysis, view the technical documentation for this indicator.

Data source: CDC, 2020a

Visit <https://www.epa.gov/roe> to see the full exhibit.

Exhibit 4. Age-adjusted coronary heart disease death rates in the U.S. by EPA Region, 1979-2018




Due to differences in the ICD system used for classifying mortalities, data from 1979-1998 should not be directly compared with data from 1999-2018 [ICD-9 codes: 410-414, 429.2 (1979-1998); ICD-10 codes: I20-I25 (1999-2018)].

Rates are age-adjusted to the 2000 U.S. standard population.

Information on the statistical significance of the trends in this exhibit is not presented here. For more information about uncertainty, variability, and statistical analysis, view the technical documentation for this indicator.

Data source: CDC, 2011, 2020a

Exhibit 5. Age-adjusted stroke death rates in the U.S. by EPA Region, 1979–2018



Due to differences in the ICD system used for classifying mortalities, data from 1979–1998 should not be directly compared with data from 1999–2018 [ICD–9 codes: 430–434, 436–438 (1979–1998); ICD–10 codes: I60–I69 (1999–2018)].

Rates are age-adjusted to the 2000 U.S. standard population.

Information on the statistical significance of the trends in this exhibit is not presented here. For more information about uncertainty, variability, and statistical analysis, view the technical documentation for this indicator.

Data source: CDC, 2011, 2020a