

Reactive nitrogen in atmospheric emission inventories

Stefan Reis¹, Robert W. Pinder², Meigen Zhang³, Gao Lijie³, Mark A. Sutton¹

srei@ceh.ac.uk

¹ Centre for Ecology & Hydrology (CEH),
Bush Estate, Penicuik, EH26 0QB, United Kingdom

² US Environmental Protection Agency, Atmospheric Modeling Division,
109 T W Alexander Drive, Research Triangle Park, NC 27711, USA

³ Institute of Atmospheric Physics, Chinese Academy of Sciences,
Beijing, People's Republic of China

1 Abstract

Excess reactive Nitrogen (Nr) has become one of the most pressing environmental problems leading to air pollution, acidification and eutrophication of ecosystems, biodiversity impacts, leaching of nitrates into groundwater and global warming. This paper investigates how current inventories cover emissions of Nr to the atmosphere in *Europe*, the *United States of America*, and *The People's Republic of China*. The focus is on anthropogenic sources, assessing the state-of-the-art of quantifying emissions of Ammonia (NH₃), Nitrogen Oxides (NO_x) and Nitrous Oxide (N₂O), the different purposes for which inventories are compiled, and to which extent current inventories meet the needs of atmospheric dispersion modelling. The paper concludes with a discussion of uncertainties involved and a brief outlook on emerging trends in the three regions investigated is conducted.

Key issues are substantial differences in the overall magnitude, but as well in the relative sectoral contribution of emissions in the inventories that have been assessed. While these can be explained by the use of different methodologies and underlying data (e.g. emission factors or activity rates), they may lead to quite different results when using the emission datasets to model ambient air quality or the deposition with atmospheric dispersion models. Hence, differences and uncertainties in emission inventories are not merely of academic interest, but

can have direct policy implications when the development of policy actions is based on these model results.

The level of uncertainty of emission estimates varies greatly between substances, regions and emission source sectors. This has implications for the direction of future research needs and indicates how existing gaps between modelled and measured concentration or deposition rates could be most efficiently addressed.

The observed current trends in emissions display decreasing NO_x emissions and only slight reductions for NH₃ in both Europe and the US. However, in China projections indicate a steep increase of both.

Keywords: nitrogen, ammonia, nitrogen oxides, nitrous oxide, emission inventories

2 Introduction

2.1 Aims and objectives

Nitrogen gas (N_2) accounts for more than 99.99% of all the nitrogen present in the atmosphere, while of the rest, again 99% is accounted for by nitrous oxide (N_2O) (Wallace and Hobbes, 2006). Other N species are thus only present in trace concentrations, but nonetheless play a vital role in atmospheric chemistry. Ammonia (NH_3) is the most abundant alkaline gas in the atmosphere and is responsible for neutralising acids formed through the oxidation of sulphur dioxide (SO_2) and nitrogen oxides (NO_x), creating ammonium (NH_4^+) salts of sulphuric and nitric acid, which become atmospheric aerosols. The only other basic gases in the atmosphere are also reduced nitrogen, such as volatile amines, though these are present in much smaller quantities. Oxidized nitrogen, mainly originating as nitric oxide (NO) and nitrogen dioxide (NO_2) plays a crucial roles both in tropospheric and stratospheric chemistry, for instance in the formation of tropospheric ozone (see Bradshaw et al., 2000; Brasseur et al., 1999).

Nitrogen is thus present in the atmosphere in a multiplicity of chemical forms, contributing both the majority (78%) of the atmosphere as N_2 and a plethora of trace N components that are fundamental to the atmosphere's chemical and radiative properties. It is convenient to distinguish these N forms into two main groups: N_2 being termed non-reactive (or 'fixed') nitrogen, with the sum of all other N forms present being termed 'reactive nitrogen' (Nr) (Galloway et al. 2003 and 2008).

The amounts of Nr in the world are of fundamental importance to society. Since the invention of the Haber-Bosch process it has become possible to synthesise huge amounts of NH_3 directly from N_2 . The key benefit has been to feed the increasing world population, which is offset by the consequent increases in Nr losses to the environment (e.g. Erisman et al., 2008). In parallel, high temperature combustion of fossil fuels oxidizes atmospheric N_2 , causing a

huge increase in NO_x emissions in the atmosphere (Lee et al., 1997). The alteration in agricultural practice and increase in fossil fuel combustion impacts human health, acidification and eutrophication of soils, biodiversity change in terrestrial ecosystems, nitrate eutrophication in freshwater and marine ecosystems and alteration of the global radiative balance.

While there is sufficient evidence to demonstrate the substantial perturbation of the global nitrogen cycle, the exact quantification of the magnitude and spatial distribution of this perturbation is presently subject to in-depth research. The NitroEurope research project (Sutton et al., 2007; <http://www.nitroeurope.eu>) is working toward deriving more precise nitrogen balances from local to regional scales. A major challenge in compiling nitrogen budgets and quantifying pools and fluxes of nitrogen in the atmosphere and biosphere is that models and measurements need to be of good and known quality to allow for a validation of results and ultimately provide the scientific understanding of processes on all relevant spatial scales. The variety of sources of different forms of nitrogen being emitted into the atmosphere, soils and water bodies, their heterogeneous distribution in space and the often high uncertainties regarding specific N fluxes creates a challenge for the validation and verification of models. At present, closing the gaps in the nitrogen budgets is the aim of major efforts on a global scale. One example for this is the European Nitrogen Assessment Report (compiled under the auspices of the ESF Programme Nitrogen in Europe and the International Nitrogen Initiative) and the , which is due to be completed by early 2011 and which aims to improve the understanding of the European N cycle and move toward closing the gaps. Emission inventories are a critical source of information in this process. However, inventories are most often compiled for compliance monitoring purposes, e.g. EMEP under the UNECE Convention on LRTAP or the United Nations Framework Convention on Climate Change (UNFCCC), using officially submitted data. On the other hand, inventories compiled by

researchers with the main aim to provide a consistent and comprehensive set of input data for modelling purposes (e.g. the EDGAR dataset) often use a bottom-up approach with information on emission factors and activity data independent of official (often not readily accessible) data.

In this paper, we focus on regional scale nitrogen emission inventories, with the aim to assess how different methods and data sources influence the resulting inventory datasets. In a second step, we analyse the main drivers for differences identified, and consider how these may affect the usefulness of datasets for atmospheric modelling. Finally, we briefly discuss emission trends from inventory datasets and scientific literature, considering the implications for our conclusions on the use of emission datasets for modelling purposes.

2.2 Scope

Table 1. Overview of the main sources and sinks of atmospheric nitrogen-containing species. N₂ has been listed for completeness, but is not discussed further, as the paper focuses on reactive N species only.

Sources	(N ₂)	NH ₃	NO _x	N ₂ O
Biogenic emissions from the terrestrial and marine biosphere		X	X	X
Decomposition of proteins and urea from animals		X		
Biomass burning and fossil fuel combustion	(X)	X	X	
Agricultural mineral Nr fertilisation and denitrification	(X)	X	X	X
Lightning			X	
Sinks				
Wet deposition (as NH ₄ ⁺ and NO ₃ ⁻)		X	X	
Dry deposition		X	X	
Biological nitrogen fixation	(X)			
Industrial nitrogen fixation (mainly Haber-Bosch process)	(X)			
Chemical breakdown in the stratosphere				X

Anthropogenic activities have a significant impact on the magnitude of N cycled and released into the atmosphere, for instance, Galloway et al. (2004) calculated a global rate of annual creation of reactive nitrogen (N_r) of 163 Tg N yr⁻¹ in the early 1990s compared to 125 Tg N yr⁻¹ around 1860.

The following Table 2 displays the total N emissions in Tg for the three regions on which this paper focuses. It illustrates both the similarities and differences between different inventories and the contribution to the atmospheric domain of global N emissions, which based on the EDGAR v4 inventory amounts to approx. 32 Tg N in the year 2005.

Table 2. Emissions in Tg nitrogen (Tg N) for the three regions analysed in this paper based on a global inventory (EDGAR v4) and national submissions (to EMEP, IPCC). In the case of the PR of China, no official national inventory was available, NH₃ emissions as estimated for 2006 by Yan et al. (2003) and other data from the IIASA GAINS China model have been used for the comparison

	EU25 ^a		USA		PR of China	
	EDGAR v4	EMEP ¹ /IPCC ²	EDGAR v4	US EPA	EDGAR v4	IIASA ³ /Yan et al. 2003 ⁴
NH ₃	4.35	3.30 ¹	2.81	3.02	8.43	5.80 ⁴
NO _x	3.51	3.49 ¹	5.48	5.61	5.59	5.20 ³
N ₂ O	0.63	0.82 ²	0.46	0.65	0.68	1.18 ³
	8.48	7.61	8.75	9.28	14.69	12.18

a) EU27, except Malta and Cyprus, omitted for consistency with figures

At the same time, the scientific understanding of many environmental effects of excess nitrogen in the atmosphere has significantly advanced in recent years. Current research into the critical loads of N deposition both for acidification and eutrophication (see Hettelingh et al., 2008) has led to the establishment of more stringent critical loads, with dynamic modelling approaches being explored to assess the timescales of ecosystem damage and recovery. In addition, the relevance of N₂O as a contributor to global warming has been acknowledged and emission control strategies no longer focus solely on the CO₂, which is reflected e.g. by a more detailed integration of N₂O sources and measures in the GAINS model (Greenhouse Gas – Air Pollution Interaction and Synergies, <http://gains.iiasa.ac.at/>) or MITERRA (Velthof et al, 2008).

Given the importance of nitrogen-containing species for air quality and climate change, the question emerges if the current inventories of NH₃, NO_x and N₂O reflect the current knowledge on emissions of these trace gases. Furthermore, the parallel but rather separated development of emission inventories under air quality related activities (such as the UNECE *Convention on Long Range Transboundary Air Pollution*) and under the climate change focus

of the *Intergovernmental Panel for Climate Change* (IPCC) and the *United Nations Framework Convention on Climate Change* (UNFCCC) has led to different accounting systems with often differing national budgets for the same trace gas. In this paper, we compare how different approaches to inventory compilation may lead to similar or quite different results and – where possible – discuss likely reasons for differences observed. Specifically, we assess whether the quality of current emission inventories is sufficient to support integrated strategies for N management, which are emerging in the US and Europe (Erisman et al., 2007). For this purpose, we here assess existing inventories with regard to their total numbers and *sectoral structure*. The *temporal* and *spatial resolution*, *speciation* and *accessibility of inventories* are not the focus of this paper, but will be addressed where necessary.

In the following sections, the current situation of emission inventories in Europe, the US and China is discussed with the focus on how emissions of NH_3 , NO_x and N_2O are estimated and allocated to source sectors. Aspects of spatial and temporal resolution, sectoral detail on emission sources and completeness of reporting are addressed for the legislative and regulatory regimes under which data are compiled. Where different inventories are compiled, a comparison and analysis of potential variations will be conducted, alongside uncertainty assessments.

For this purpose, the following inventories and data sources have been analysed in detail:

- National submissions of NO_x and NH_3 emission data to the EMEP programme
- National submissions of N_2O emission data to the UNFCCC/IPCC
- The US National Emissions Inventory (NEI) dataset for NO_x and NH_3
- Inventory of U.S. Greenhouse Gas Emissions and Sinks for N_2O

- The EDGAR global emission inventory dataset for NO_x, NH₃ and N₂O (version 4 representing the most up-to-date emissions; EDGAR Fast Track 32 and EDGAR HYDE for trends and for comparison purposes)
- Literature data in particular for the PR of China and for global/regional comparisons.

The datasets listed are openly accessible, in most cases directly online. The EDGAR v4 inventory has only been partly published at this stage (greenhouse gases). However, the authors had access to a preliminary version of the dataset for air pollutants, which will be published in the near future, subject to error corrections to which the comparisons made in this paper may contribute (<http://edgar.jrc.ec.europa.eu/>).

On a global scale, the need for nitrogen management has been formulated in the 2004 Nanjing Declaration (Erisman 2004, UNEP 2004), which was presented to the *United Nations Environment Programme* (UNEP) in Nanjing, China on October 16, 2004. With the aim to optimise nitrogen management in food and energy production on a local, regional and global scale, the declaration calls upon the national governments for instance to

- support the further assessment of the nitrogen cycle,
- focus efforts on increasing the efficiency and effectiveness of agricultural production and energy use, while decreasing the adverse effects of reactive nitrogen, and
- take action to enhance the availability of reactive nitrogen as food, fibre and other basic needs in regions of nitrogen deficiency and avoid nitrogen pollution.

For this last step, three key approaches are identified:

1. A code of good agricultural, forestry and aquacultural practise.
2. Strategies for sustainable energy use to prevent the formation of nitrogen oxides in fossil fuel combustion.

3. Application of emission control technologies.

With regard to the requirements for an integrated nitrogen management and the need to fulfill different obligations under e.g. the UNECE CLRTAP or the UNFCCC, we investigate here if the current methods to account for the emissions of different nitrogen species in emission inventories provide a sufficiently robust data basis for such in depth integrated assessments.

3 European emission inventories of reactive nitrogen species

3.1 Overview

In Europe, emissions of ammonia and nitrogen oxides are covered by several regulatory regimes, both under the UNECE CLRTAP and directives of the European Commission. Member states of the European Union and parties to the protocols under the CLRTAP are subject to mandatory emission reporting. For the UNECE, the EMEP Centre on Emission Inventories and Projections (CEIP) at the Umweltbundesamt Vienna, Austria, hosts inventory datasets (both official submissions of signatories to the different protocols of the CLRTAP and emissions for modelling purposes) in an online-accessible database (<http://www.emep-emissions.at/emission-data-webdab/>)

Nitrous oxide on the other hand is not covered by the CLRTAP, but is required to be reported under the UNFCCC by ANNEX I countries. The UNFCCC GHG inventory submissions can be accessed http://unfccc.int/ghg_emissions_data/items/3800.php online.

In addition to these inventories which are generated based on obligatory reporting of national emissions, the EDGAR database provided global annual emissions per country and on a 1×1 degree grid for 1990 and 1995 for direct greenhouse gases CO₂, CH₄, N₂O and HFCs, PFCs and SF₆ and the precursor gases CO, NO_x, NMVOC and SO₂. Similar inventories have been compiled for acidifying gases, NH₃, NO_x and SO₂ and Ozone Depleting Gases (EDGAR 2008). For this paper, the new EDGAR dataset version 4 (v4 in the following text) is used,

which has recently partly (GHGs) been officially released and of which a preliminary version was made available to the authors by the EDGAR team (<http://edgar.jrc.ec.europa.eu/>). EDGAR v4 (EDGAR, 2008) provides emissions for all relevant air pollutants and GHGs with an improved spatial resolution and until 2005.

3.2 Ammonia

In the case of NH_3 , the vast majority of emissions from the 27 member states of the European Union (EU27) originate from agriculture (93%), with some small contributions from waste management (2.5%), industrial production processes (2%) and road transport (1.8%) (EMEP, 2008). This sectoral distribution is valid for most countries, with slight difference depending on the state of the art of agricultural production and, for instance, livestock intensity. For ammonia, a large number of non-agricultural sources contribute a small amount of emission (Sutton et al., 2000). Because the individual contributions are small for these sources (e.g. wild animals, direct emissions from humans, sewage management) many countries do not report emissions for all these terms. For example, detailed analysis of these non-agricultural emissions for the UK showed that they contribute around 15% of total ammonia emissions (Sutton et al., 2000; Dragosits et al., 2008). This is double the share noted above for the EU27 as a whole, which clearly indicates that a more comprehensive reporting of NH_3 emissions is needed.

Fig. 1 illustrates that differences between the EMEP dataset and data used for model calculations with the RAINS model by IIASA are marginal for most countries. This was anticipated, as the EMEP emissions displayed represent official submissions by countries, which, in ultimately, form the basis for the IIASA data through a validation process by extensive bilateral consultations with country experts, often the same experts preparing the inventories reported to EMEP.




Fig. 1. Comparison of EU25 (not showing Malta and Cyprus) emissions of ammonia (expressed in Gg N) reported to EMEP and compiled by IIASA and the EDGAR v4 database for the year 2005. The slight differences between IIASA estimates and EMEP figures arise most likely from bilateral consultations with country experts, which led to corrections in agricultural emissions that had not (yet) been reflected in the EMEP inventories by recalculations of the year 2000 emissions submitted.

For some countries, however, the IIASA data are slightly different, with overall emissions for the EU27 being 4.4% below the EMEP figures. The reason for this small difference in emissions that have the same underlying data sources is most likely revised animal numbers or more detailed emission calculations that have been available to the experts during the consultations, but have not yet been used to submit recalculated inventory figures to EMEP. The comparison with the EDGAR v4 dataset (EDGAR, 2008) shows that EDGAR emissions are (consistently) higher for the bulk of EU27 countries (32%). As a major difference, the EDGAR emissions for agricultural sources are significantly higher than those reported to EMEP, with emissions from agricultural soils being most likely the main contributor to the difference observed. A more detailed analysis is not straightforward due the degree of

completeness of EMEP emissions reported based on the current reporting format (Nomenclature for Reporting, NFR08), which distinguishes in sufficient detail emissions within the agricultural sector (currently available for 10 countries). An in-depth assessment is thus conducted exemplary for the case of the UK in Section 6.2.2.

For the European region, including the EU27, the accession countries (Turkey, FYR of Macedonia and Croatia), as well as Norway and Switzerland, the EMEP inventory amounts to 4.55 Tg of NH_3 for the year 2005 (3.3 Tg N). This is comparable with an estimate of 4.1 Tg NH_3 for the whole of Europe made by Bouwman et al. (1997) for the year 1990 and an estimate of 5.3 Tg NH_3 (4.3 Tg N) by EDGAR (2008) for the same set of European countries. Galloway et al. (2004) only give a combined figure for Europe and the former Soviet Union (FSU), with atmospheric emissions of ammonia calculated at 8 Tg N yr^{-1} , with FSU emissions in 1990 at 3.4 Tg N yr^{-1} according to the EDGAR dataset.

3.3 Nitrogen Oxides

Nitrogen oxides have been the focus of significant emission control activities in recent decades, both for stationary sources (mainly large combustion plants) and mobile sources (especially road transport). In general it should be anticipated that NO_x emission figures are less uncertain than those of NH_3 or N_2O and the initial comparison between EMEP and EDGAR datasets for 2005 confirms this with only slight differences for individual countries (see Fig. 2). Total NO_x emissions (expressed as $\text{NO}+\text{NO}_2$) for Europe are estimated at 11.5 Tg by both EDGAR v4 and EMEP (EDGAR, 2008; EMEP, 2008) for the year 2005 (3.5 Tg N).




Fig. 2. Comparison of EU27 (not showing Malta and Cyprus) emissions of NO_x (expressed as Gg N) reported to EMEP (EMEP, 2008) and presented in the EDGAR database (EDGAR, 2008) for the year 2005

This comparison indicates that NO_x emissions are undoubtedly better understood than for instance NH₃, in general. National totals do not display large variations between inventories, however, sectoral differences can be significant for individual countries. Issues such as an overall lack of measurement programmes, for instance for new vehicle technologies in road transport, and the uncertainties in the effects of decentralised power generation in a liberalised energy market on power plant emissions are likely to have an effect on the quality of NO_x inventory datasets also in the future. For an in-depth assessment of e.g. sectoral differences between inventories, up-to-date and documented national emission factors based on measurements for different technologies would be vital.

3.4 Nitrous Oxide

Since N₂O emissions are not reported under the CLRTAP, but subject to reporting obligations to the UNFCCC for ANNEX 1 countries, the comparison is made between data collected under the UNFCCC and the EDGAR v4 inventory (Fig. 3). For most countries, figures in both inventories are quite similar, with EDGAR showing an overall lower emission for the EU27 of about 24%. For comparison, figures from the IIASA GAINS model are displayed in Fig. 3 as well.




Fig. 3. Comparison of EU27 (not showing Malta and Cyprus) emissions of N₂O (expressed as Gg N) reported to UNFCCC (without LULUCF, UNFCCC, 2008) and presented in the EDGAR v4 database (EDGAR, 2008) for the year 2005

While EDGAR shows higher emissions for France (23%), most other figures are substantially lower than UNFCCC data, for instance Finland (-45%), Italy (-60%) or Sweden (-47%). At this stage, it is difficult to assess fully the uncertainty in either dataset, but it should be stated that recent findings of Skiba et al. (2001) provide a methodology for the calculation of N₂O emissions from soils, one of the main sources of N₂O,

which results in higher emissions than the current UNFCCC established methodology. Within the EU27, 48% of N₂O emissions in the EDGAR v4 inventory stem from agricultural sources (34.2% from agricultural soils, 13.9% from manure management and 0.1% from agricultural waste burning). Shares of emissions from agricultural soils in individual countries vary significantly, between 9.6% (Greece) and 67.6% (Denmark). Fig. 4 highlights the variation of relative contributions of agricultural soils and manure management to N₂O emissions in the EU27 countries. A third contributing source to N₂O emissions in the EU27 is the chemical industry sector, contributing about 28% of total N₂O emissions in 2005. It is likely that emissions from the chemical industry are well understood and hence emission factors and activity rates can be assumed to be less uncertain than those from agricultural soils or manure management. Hence, the differences between UNFCCC and EDGAR data most likely mainly arise from the application of different emission factors (or activity rates) in these two sectors.




Fig. 4. Relative contributions of N₂O emissions (in Gg N₂O) from agricultural sources soils and manure management in the EU27 countries (not displaying Cyprus and Malta) according to the EDGAR v4 inventory (EDGAR, 2008) and compared to major agricultural sources as reported to UNFCCC (Agricultural Soils, NRF 4D; Manure Management, NRF 4B) (UNFCCC, 2009)

Fig. 4 illustrates the difference between the main sources of N₂O emissions with a focus on emissions from agricultural sources. There is a clear indication that emissions reported to UNFCCC under CRF (Common Reporting Format) categories 4B (Manure Management) and 4D (Agricultural Soils) are substantially higher than those estimated in EDGAR v4 under the same headings.

However, even in those cases where both inventories provide similar figures, a more thorough investigation of emissions from agricultural soils may be required in the light of the findings of Skiba et al. (2001), Crutzen et al. (2007) and Mosier et al. (1998), which indicate a potential underestimation of N₂O emissions from soils in current inventories. The increasing demand for biofuels could even lead to a larger underestimation, unless future emission factors take these findings into account.

4 Emission inventories of reactive nitrogen species in the United States of America

The U.S. Environmental Protection Agency is charged with developing the National Emission Inventory (NEI, US EPA, 2009a) in support of the Clean Air Act and subsequent amendments. The NEI includes an accounting of pollutants that impact air quality, including NO_x and NH_3 . These efforts have recently been reviewed in an assessment report by NARSTO (2005). In addition, the EPA prepares an estimate of N_2O emissions in the U.S. Inventory of Greenhouse Gas Emissions and Sinks (US EPA, 2009b), in accordance with the UNFCCC.

4.1 Ammonia

The NEI estimates that more than 80% of total USA ammonia emissions are from livestock manure management and application of chemical fertilizers. The next largest source is ammonia from vehicles equipped with catalytic converters, which comprises approximately 7% of the inventory. While the NEI and EDGAR database have similar total agricultural and vehicle emissions, 14% of the EDGAR NH_3 emissions are from industrial combustion, while this source is less than 1% in the NEI.

Because of the operational challenges in measuring ammonia emissions and a lack of detailed animal husbandry practices data, ammonia emission estimates have high uncertainty. Independent efforts to quantify the seasonal variability have shown agreement for winter and summer emissions, but differ for the spring and fall (Gilliland et al., 2006; Pinder et al., 2006; Henze et al., 2008). Because atmospheric agricultural emissions are rarely regulated, the trend in emissions is expected to be proportional to the increase in livestock population and acres under cultivation. Both are expected to increase in coming years (USDA, 2007).

4.2 Nitrogen Oxides

The NEI estimates that the largest contributors of US NO_x emissions include on-road vehicles (32%), off-road vehicles (30%), such as ships, aircraft, and construction equipment, and electricity and industrial power generation (27%). Emissions from on-road and off-road vehicles are calculated using the National Mobile Inventory Model (NMIM) (<http://www.epa.gov/otaq/nmim.htm>). While not used in this study, a more advanced mobile source emissions model, MOVES (<http://www.epa.gov/otaq/models/moves/>), is currently under development and a draft version is available for public use. Important improvements include more detailed inspection and maintenance data and better representation of extended idling emissions.

Despite increased diesel and gasoline consumption, NO_x emissions from on-road sources are estimated to have decreased by on average 5% per year since 2002 due to the introduction of stricter emission standards. Parrish (2006) used ambient data to show that total on-road emissions may have increased from 1990-2000; however, in a multi-city, multi-year study, Bishop and Stedman (2008) have demonstrated that per-car emission factors have decreased from 2000 to 2006. The NEI reports little year-to-year change in off-road vehicle emissions. Many electricity and industrial power generation facilities have been subject to several recent regulated emission reductions including the NO_x Budget Trading Program. Because most of these facilities are equipped with Continuous Emission Monitors, the emission magnitude and trend is well quantified. Since 1999, this sector has reduced emissions by an average of 6.4% per year. The combined effect of the reductions in mobile and stationary source NO_x emissions has been observed in the surface concentration monitoring data (Godowitch et al., 2008) and from space-based remote sensing methods (Kim et al., 2006; Kim et al., 2009).




Fig. 5. Comparison of US EPA emissions of N₂O, NH₃, and NO_x with the EDGAR database for the year 2005 (USEPA, 2009ab; EDGAR, 2008).

In Fig. 5, EDGAR v4 and US EPA NO_x emission data are aggregated by sector to make them directly comparable. Where stationary combustion sources are concerned, EDGAR has markedly larger emissions. The NEI power generation emission rates are likely more accurate since they are measured at the electricity generating units as part of the Continuous Emission Monitoring System. Road transport emissions are similar in both datasets, but interestingly non-road mobile sources (which include off-road vehicles, rail, air and shipping) are

significantly larger in the US EPA dataset. The NEI partially includes international shipping emissions near the US coast, which have been omitted from the EDGAR inventory and have been estimated in EDGAR FT32 at about 1.9 Tg for the US in the year 2000. Another possible explanation is a difference in activity rates and/or emission factors for domestic air transport. From 2000 to 2005, the trends in these two databases also differ. For on-road sources, EDGAR estimates 6% yr⁻¹ reduction (NEI: 4% yr⁻¹), but for power generation and industrial sources, the EDGAR inventory has little trend (NEI: 6% yr⁻¹ reduction).

4.3 Nitrous Oxide

Total U.S. N₂O emissions for 2005 are estimated to be 1.02 Tg N₂O (US EPA, 2009b). The largest source of US N₂O emissions is agricultural soils (67%), followed by fossil fuel combustion in vehicles (12%), industrial processes (8%), fossil fuel combustion for electricity generation (5%) and livestock manure management (4%). Enhancements of N₂O emissions from agricultural soils include practices such as fertilization, application of livestock manure, grazing animals on pasture or feedlots, and cultivation of N-fixing crops. The estimated trend in N₂O emissions for the USA is a gradual reduction of approximately 1% per year since 2000. This is due to a 5% reduction per year in the emissions from vehicles and industrial processes – the other sectors are estimated to have largely remained constant over this period. EDGAR v4 emissions are lower than those reported by EPA. This is almost entirely due to lower emissions from agricultural soils in the EDGAR v4 inventory.

The physical and chemical processes that drive emissions of NH₃, NO_x, and N₂O are often intertwined. This is also true for the human activities that cause these emissions, such as fuel combustion in motor vehicles and agricultural land under cultivation. However, the GHG emission inventory and the NEI do not always use the same models and data sources. For example, direct fertilized crop emissions of N₂O are estimated using the DAYCENT model (Del Grosso et al., 2001), while the temporal pattern of NH₃ emissions are estimated using an

emission factor approach (Goebe et al., 2003), and NO emissions are calculated using the BEIS model (<http://www.epa.gov/asmdnerl/biogen.html>). Because the interactions between these processes are not explicitly considered, these shortcomings impede efforts to devise strategies that simultaneously mitigate climate change and improve air quality.

5 Emission inventories of reactive nitrogen species in the People's Republic of China

The analysis of the emission situation in the PR of China is comparatively more difficult than it is the case for Europe or the United States. Reasons for this are, among others, the immense growth rates of the economy and thus fast developing emission sources and the lack of an official national inventory programme, respectively no reporting of emissions to international organisations. Fig. 6 displays the emission estimates for the year 2005, distinguishing the main source sectors. NH_3 and N_2O emissions are here dominated by agricultural sources (mainly agricultural soils and manure management), while for NO_x , the main difference to Europe and the US is the large share of stationary combustion relative to mobile sources. In the following sections, recent studies are discussed for each of the substances.




Fig. 6. Emissions of NH_3 , NO_x and N_2O (expressed in Tg N) for the PR of China in 2005 according to the EDGAR v4 inventory (EDGAR, 2008) for the main emitting source sectors. N_2O emissions are increased by a factor of 10 for ease of comparison.

5.1 Ammonia

Only a few studies on ammonia emissions in China are available (e.g. Zhao et al., 1994; Yan et al., 2003). With regard to total NH_3 emissions in China, the major contribution comes from N-fertilizer application and livestock, representing in the 1990's about 52% and 41% of total emissions, respectively (Klimont, 2001a). Other sources of ammonia emissions include biomass burning, natural ecosystems, crops and oceans, humans (breath, sweat, excretion) and fossil fuel combustion.

The basic methodology applied to derive these emissions relies on the approach used in Europe (Klaassen, 1994; Klimont, 2001b), and as far as available takes into account information about China-specific characteristics. Klimont (2001a) estimated the total ammonia emissions in China at 9.7 and 11.7 Tg in 1990 and 1995. Spatial distribution of emissions was also presented at a $1 \times 1^\circ$ grid resolution. In 1995 the highest ammonia emission density which exceeds 100 Gg NH_3 per grid, is observed in *Jiangsu* and *Henan* provinces. This corresponds well with the large population of pigs in these regions as well as high cattle density in *Henan* province. Using the IPCC approach, NH_3 emission from synthetic fertilizer and manure application in 1990 was estimated to be 1.65 Tg N by Li Yu'e et al. (2000). Yan et al. (2003) quantified the use of urea and ammonium bicarbonate and the cultivation of rice leading to a high average ammonia loss rate from chemical N fertilizer in East, Southeast and South Asia, and the total emission was estimated to be 5.8 Tg N for the area of China.

Due to anticipated increases in synthetic fertilizer application rates and per-capita meat and dairy consumption, future ammonia emissions are expected to continue to rise.

industrial installations were the largest individual source group, contributing approx. 42% of total emissions (5.0 Tg). From 1995 to today, some studies (e.g., Aardenne et al., 1999; Streets et al., 2000) estimated that NO_x emissions in China will continue to grow rapidly. However, other researches' results using latest available statistical data (e.g., Tian et al., 2001) indicate that NO_x emissions began to decrease somewhat, with total emissions in China amounting to 11.3 Tg NO_x (as NO₂) in 1995, 12.0 Tg (1996), 11.7 Tg (1997) and 11.2 Tg (1998). In the analysis by Tian et al., this is explained on the basis of changing energy management in China. According to Streets et al. (2003) Chinese emissions in the year 2000 have slightly increased again (11.3 Tg). There is a clear indication that the total NO_x emissions in China are not growing as rapidly as some research had previously projected (e.g., Aardenne et al., 1999). Satellite observations of the tropospheric NO₂ column density over Eastern China have increased considerably during 1996 to 2004 (Richter et al., 2005; van der A et al., 2006), suggesting large emission increases over that period.

5.3 Nitrous Oxide

The UNDP/GEF ECPINC Project (*Enabling China to Prepare Its Initial National Communication*, <http://www.ccchina.gov.cn/en/NewsInfo.asp?NewsId=5392>) has been implemented to support China in fulfilling its commitments under the UNFCCC to communicate to the Conference of Parties to the Convention : (1) a national inventory of emissions and sinks of greenhouse gases , (2) a general description of steps taken or envisaged by China to implement the Convention , and (3) any other information China considers relevant and suitable for inclusion in its Communication.

A fair amount of research has been conducted on an N₂O emissions inventory for China. Zheng et al. (2004) have shown that up to most (up to 75%) of cropland N₂O emissions are

direct emissions: immediately from fertilized top-soil rather than denitrification of nitrate leached into sub-surface and groundwater. Zheng et al (2004) collected 54 direct N_2O emission factors (EF_d 's) obtained from 12 sites of Chinese croplands and found that of these 60% are underestimated by 29% and 30% are overestimated by 50% due to observation shortages. The biases of EF_d s are corrected and their uncertainties are re-estimated. Of the 31 site-scale EF_d s, 42% are lower by 58% and 26% are higher by 143% than the IPCC default values. Periodically wetting/drying the fields or doubling nitrogen fertilizers may double or even triple an EF_d . The direct N_2O emissions from Chinese croplands are estimated at $275 \text{ Gg N}_2\text{O-N yr}^{-1}$ in the 1990s, of which $\sim 20\%$ is due to vegetable cultivation. The great uncertainty of this estimate, -79% to 135% , is overwhelmingly due to the huge uncertainty in estimating EF_d s ($-78 \pm 15\%$ to $129 \pm 62\%$). The direct N_2O emission intensities significantly depend upon the economic situation of the region, implying a larger potential emission in the future.

However, agricultural activity is not the only source of N_2O emissions in China. As the statistical results by Li Yu'e et al. (2000), total N_2O emissions from fuel combustion amounted to $58.22 \text{ Gg N yr}^{-1}$ in 1990 in China. Within the fuel combustion sector, energy industries, manufacturing industries and residential areas were the main sources of N_2O . Among fossil fuels, hard coal was the main contributor to N_2O emissions. The industrial process sector was a less critical source of N_2O emissions, with the total N_2O emissions from this source group amounting to $0.41\text{--}0.90 \text{ Gg N yr}^{-1}$. The total emission of N_2O as a result of fertilizer application was $342.5 \text{ Gg N yr}^{-1}$ in 1990, being the most important contributor to N_2O emission from agricultural soils.

While both Li and Xing's estimations did not consider permanent croplands, Lu et al (2006) established an empirical model to develop a spatial inventory at the $10 \times 10 \text{ km}$ scale of direct

N₂O emissions from agriculture in China, in which both emission factor and background emission for N₂O were adjusted for precipitation. As a result, the total annual fertilizer-induced N₂O emission was estimated to be 198.89 Gg N₂O-N in 1997 and background emissions of N₂O from agriculture was estimated to be 92.78 Gg N₂O-N and the annual N₂O emission totalled 291.67 Gg N₂O-N.

All N₂O emission measurements are subject to significant uncertainty due to their great temporal and spatial variations of cropland fluxes.

6 Evaluation of emission inventories for reactive nitrogen

6.1 Uncertainty assessment in general

Table 3. Indication of uncertainty estimate for greenhouse gases.
Source: Olivier et al., 1999

Emission source category	Activity data	Emission Factor			Total Emissions		
		CO ₂	CH ₄	N ₂ O	CO ₂	CH ₄	N ₂ O
<i>Fossil fuel use</i>							
Fossil fuel combustion	S	S	M	M	S	M	M
Fossil fuel production	S	M	M	-	M	M	-
<i>Industry/solvent use</i>							
Iron & steel production	S	-	S	-	-	S	-
Non-ferrous production	S	-	S	-	-	S	-
Chemicals production	S	-	S	L	-	S	M
Cement production	S	S	-	-	S	-	-
Solvent use	M	-	-	-	-	-	-
Miscellaneous	V	-	-	-	-	-	-
<i>Landuse/waste treatment</i>							
Agriculture	S	-	L	L	-	L	L
Animals (excreta/ruminants)	S	-	M	L	-	M	L
Biomass burning	L	S	M	L	L	L	L
Landfills	L	-	M	-	-	L	-
Agricultural waste burning	L	-	L	L	-	L	L
Uncontrolled waste burning	L	-	-	-	-	-	-
All sources	-	-	-	-	S	M	L

S = small (10%); M = medium (50%); L = large (100%); V = very large (>100%) "-" not applicable/negligible

For all emission inventories portrayed here, the assessment of uncertainties is a key aspect. For national inventories, issues of compliance with reduction targets and emission ceilings are relevant, while in general the quantification of uncertainties of emissions as input data for

atmospheric dispersion models is of importance. Aspects of completeness regarding the total amount of emissions accounted for are as important as the spatial distribution, chemical composition and temporal patterns of emission occurrences.

Within the context of emission inventories compiled under the CLRTAP, country submissions are accompanied by so-called informative inventory reports (IIRs, <http://www.emep.int/emis2007/reportinginstructions.html>), covering aspects of completeness, an analysis of key sources and uncertainties. In addition to this, regular centralised reviews are conducted for selected countries, with the aim to improve the quality and accuracy of reported emission data.

Under the UNFCCC, emission reporting is guided by a document called “Good Practice Guidance and Uncertainty Management in National Greenhouse Gas Inventories” (<http://www.ipcc-nggip.iges.or.jp/public/gp/english/>, see as well IPCC 1997, 2000) in order to *“...to assist countries in producing inventories that are neither over nor underestimates so far as can be judged, and in which uncertainties are reduced as far as practicable.”*

In addition to these institutionalised approaches to uncertainty assessment, studies such as conducted by Winiwarter and Rypdal (2001) and Rypdal and Winiwarter (2001) provide a methodological analysis of uncertainties for specific trace gases, respectively individual countries. Olivier et al. (1999) have discussed uncertainties of GHG emissions on a sectoral level, distinguishing between uncertainties in activity datasets, emission factors and the resulting total emissions (see Table 3).

Olivier et al. (1999) conclude that emissions from fossil fuel production and combustion are generally well understood and prone to small to medium uncertainties only. This can be said as well for industrial production processes or solvent use. In contrast, emissions related to agricultural land-use are viewed as significantly more uncertain. In a similar way, emission estimates of N₂O and CH₄ are generally less robust, because CO₂ is emitted from combustion

processes mainly can be calculated more accurately than emissions based on soil microbial processes for instance. These findings have implications as well for the discussion of the different substances in the inventories analysed in this paper, where NO_x emissions can be seen comparatively robust. Even though the uncertainty associated with NO_x emissions is approx. 4 times higher than for CO₂, the latter are directly linear to fuel input in combustion sources and EFs are not influenced by processes and hence easier to determine. On the other hand, NH₃ and even more so, N₂O emissions are likely to be more uncertain because their main sources are related to agriculture.


For individual inventories, e.g. the UK NAEI, uncertainty assessments are provided by inventory compilers (http://www.naei.org.uk/emissions/emissions_2007.php?action=notes1). For NO_x, the NAEI assumes $\pm 7\%$, for NH₃ $\pm 20\%$, while for N₂O no individual assessment is made, but an overall uncertainty of $\pm 15\%$ is estimated across the 6 greenhouse gases overall.

6.2 Inventory comparison – analysing differences and similarities

6.2.1 General observations for the situation in Europe

A full and detailed intercomparison of the inventories on the most detailed sectoral level for all countries/regions is beyond the scope of this paper. However, some similarities and differences are worth noting. In the case of ammonia, both the EMEP and the IIASA figures are quite similar, with some countries showing different emission levels, most likely due to recalculations and re-assessments of e.g. animal numbers that have been emerging in the course of bilateral consultations in the preparation of the IIASA dataset and which have not (yet) been incorporated in recalculations of the data reported to EMEP. With regard to NO_x, the differences between EMEP and EDGAR figures are substantial for some countries and remarkably different in total (EDGAR v4 32% higher than EMEP). Some of the potential sources of these discrepancies have been highlighted in *section 3.3*, in particular the different sectoral allocations including some emission sources that are not reported under EMEP in the

EDGAR dataset. In addition to that, a likely reason for these large differences are assumptions regarding underlying emission factors for some of the largest contributing sources, e.g. due to the estimated share of power plants equipped with efficient NO_x control equipment, or emission factors and activity rates for road transport sources.






Fig. 7. Sectoral comparison of emissions for the year 2005 between EMEP (top; EMEP, 2008) and EDGAR v4 (bottom; EDGAR, 2008). It should be noted, that the EMEP sectoral split is based on only 10 countries reporting according to the latest sectoral structure (NFR 08, Level 1), which is best comparable with the EDGAR v4 sector structure.

Fig. 7 displays a direct comparison of NH_3 and NO_x emissions in the EMEP and the EDGAR v4 inventories by sector for the year 2005. The most recent change in EMEP reporting requirements, adopting a new reporting format termed NFR 08 (Nomenclature for Reporting) which is – on an aggregated level – better suited for sectoral analysis than its predecessors (e.g. NFR 01, NFR 02) which were based on the UNFCCC CRF format and in some areas lacking a sufficient level of detail. With regard to NH_3 , agricultural emissions show a similar pattern in both inventories overall and are dominating emissions for this substance. The situation for NO_x emissions is more diverse and shows a distinct difference for stationary combustion sources, which contribute more than 55% to EDGAR v4 NO_x emissions, but only account for about 42% of EMEP NO_x emissions. A full comparison of individual sectors by country on a European scale would be worthwhile once all countries are reporting emissions based on the new NFR 08 sectoral structure. As this is not feasible at this stage and within the

scope of this paper, an exemplary analysis is conducted for the United Kingdom in the following section.

6.2.2 Detailed analysis for the United Kingdom

For a thorough investigation of the differences between inventories, it is essential to have access to a very detailed sectoral split, and if feasible, even the emission factors and activity rates that have been used to compile the inventories. In this context, the structure initially applied in the EMEP inventories labelled SNAP (Selected Nomenclature for Air Pollution, EMEP) provided emission data with a detailed level 3 split allowed for a comparison down to process and fuel level. However, not many countries provided the obligatory information in this detailed split and hence inventories were subject to different levels of detail and substantial gaps. The new format for reporting under EMEP has been termed Nomenclature for Reporting (NFR, current version is NFR 08). While NFR has been closely aligned with the UNFCCC common reporting format (CRF) and thus been mainly driven by fossil fuel combustion sources, it provided less detail and no distinctions had been made for instance with regard to the type of fuel used in power plants or road transport modes. The recently adopted NFR 08 better accounts for non-combustion sources and already on the most aggregated level (NFR 08, Level 1) provides a useful split into the main sectors (for details on NFR 08, see <http://www.unece.org/env/documents/2008/EB/EB/ece.eb.air.2008.4.e.pdf>). The sectoral split used in EDGAR v4 reflects the bottom-up character of the inventory. It is not always directly comparable to NFR sectors, but in most cases, an equivalent source sector allocation – on a more aggregate level - can be found.

Table 4 provides a detailed comparison of source sectors split into categories for which a cross-comparison between the UK NAEI and EDGAR v4 could be sensibly conducted for NO_x, NH₃ and N₂O. As some of the individual source sectors in NAEI and EDGAR do not

fully match, the following discussion will focus on the subtotals for sector groups rather than on individual figures.

The figures for stationary combustion are quite similar overall, but NO_x emissions in EDGAR v4 are 17% lower, N₂O from this source group is less than a third than those in the NAEI. For road transport, the picture is similar (17% lower NO_x emissions, 87% lower N₂O emissions), while NH₃ from road transport is 47% higher in EDGAR v4. NO_x emissions from other mobile sources and machinery are about 32% lower in EDGAR v4. These combustion sources, both stationary and mobile, are assumed to be best understood with low uncertainties typically attributed to NO_x emission factors in general. Thus, the magnitude of differences in these sectors between NAEI and EDGAR v4, however, is larger than anticipated taking into account the uncertainty ranges assumed for the UK NAEI (see section 6.1).

On the other hand, total NH₃ emissions from the sector manure management are quite similar (~6% lower in EDGAR v4), yet some significant differences between the emissions from individual animal types can be observed. EDGAR v4 has substantially higher emissions of N₂O from manure management (about 6 times higher than NAEI), but less than one third of direct emissions from agricultural soils. This may hint at a difference in allocation between these sub-sectors. In addition to that, NH₃ emissions directly originating from agricultural soils are 5.3 times higher in EDGAR v4.

Finally, NAEI lists a few sectors that are not subject to reporting obligations under EMEP, while EDGAR v4 – as an inventory primarily compiled for modelling purposes – does not make this distinction.

In summary, the difference between country total emissions between both inventories is within the uncertainty margins expected for N₂O at 2.6% ($\pm 15\%$ uncertainty range for NAEI, see section 6.1), however larger for NO_x emissions with EDGAR v4 being 20.9% below NAEI, (NAEI assuming a $\pm 7\%$ uncertainty range). NH₃ emissions in EDGAR v4 are 36.9%

higher than in the NAEI (uncertainty range for NAEI assumed to be $(\pm 20\%)$). However, for all three substances, sectoral allocation are quite different between inventories, which has direct implications for the use of the emissions for atmospheric modelling, e.g. the temporal profiles and spatial distribution of emissions in different sectors. The current structure of the inventories such as NAEI and hence EMEP - which are compiled primarily for reporting and compliance monitoring purposes - is not always detailed enough for modellers to fully assess the quality of the inventory and supporting datasets are often not accessible or documented well enough.

Table 4. Detailed comparison between source sectors in the UK National Atmospheric Emissions Inventory (NAEI, 2008) and EDGAR v4 (EDGAR, 2008) for the year 2005.

NFR code	NFR Name	UK NAEI			EDGAR v4		
		NO _x	NH ₃	N ₂ O	NO _x	NH ₃	N ₂ O
1A1a	Public Electricity and Heat Production	372.60	0.70	3.48	410.14	0.57	2.65
1A1b	Petroleum Refining	30.70		0.39	3.20		0.01
1A1c	Combustion in Manufacture of Solid Fuels and Other Energy Industries	58.80		1.28			
1A2a	Combustion in Iron and Steel Manuf. Industry	19.50		0.33	12.19		
1A2f	Combustion in Other Manufacturing Industry	232.80	0.40	4.00	165.73	0.17	0.36
	Subtotal stationary combustion	714.40	1.10	9.48	591.26	0.75	3.02
1A3a	Civil Aviation - Domestic Take-Off and Landing	1.80		0.02			
1A3a(ii)	Civil Aviation - Domestic Cruise	7.10		0.05			
	Subtotal aviation	8.80		0.07			
1A3bi	Road Transport - Passenger Cars	215.10	9.20	14.13			
1A3bii	Road Transport - Light Duty Vehicles	58.50	0.20	1.19	457.04	13.98	2.14
1A3biii	Road Transport - Heavy Duty Vehicles	274.00	0.10	1.09			
1A3biv	Road Transport - Mopeds & Motorcycles	1.30		0.01			
	Subtotal road transport	548.90	9.50	16.42	457.04	13.98	2.14
1A3c	Railways - Mobile Sources	36.90		0.80			
1A3dii	National Navigation (including Inland Waterways and Maritime Activities)	92.10		0.10			
1A3eii	Other Off-Road Mobile Sources and Machinery	5.60		0.19	91.06	0.03	0.05
	Subtotal other mobile sources	134.60		1.09	91.06	0.03	0.05
1A4a	Commercial/Institutional Combustion Plants	20.90		0.08	110.14	0.77	0.51
1A4bi	Residential Combustion Plants	108.30	1.50	0.37			
	Subtotal residential/commercial combustion	129.20	1.50	0.45	110.14	0.77	0.51
1A4bii	Household and Gardening (Mobile Machinery)	0.90		0.02			
1A4ci	Stationary Combustion Plants Agriculture/Forestry/Fishing)	0.70		0.01			
1A4cii	Agricultural/Forestry Off-Road Vehicles and Other	52.10		1.59			
1A5b	Other Mobile Sources (including Military)	22.20		0.08			
1B1b	Fugitive Emissions from Fuels - Solid Fuels/Transformation	0.20	0.10	0.00			
1B2ai	Fugitive Emissions from Fuels, Oil - Exploration, Production, Transport	0.70	0.10	0.00			
1B2c	Oil and Natural Gas/Venting and Flaring	2.20		0.14			
2A7	Other Production		0.50				
2B2	Nitric Acid Production	0.60		6.52			
2B3	Adipic Acid Production			2.50			
2B5	Other Chemical Industry Processes	0.40	4.00			0.67	66.14
2C	Metal Production	1.60		0.03			
2D1	Pulp and Paper Production						
2D2	Food and Drink Production		0.90				
3D	Other Solvent Use (including products containing HMs and POPs)		1.20				

NFR code	NFR Name	UK NAEI			EDGAR v4		
		NO _x	NH ₃	N ₂ O	NO _x	NH ₃	N ₂ O
	Subtotal miscellaneous sources	157.50	6.80	12.59		0.67	66.14
4B12	Solid Storage and Drylot			0.15			
4B13	Manure Management - Other		17.90	3.80			
4B1a	Manure Management - Dairy		86.00			52.20	5.72
4B1b	Manure Management - Non-Dairy		63.60			75.92	8.10
4B3	Manure Management - Sheep		12.10			23.49	7.33
4B6	Manure Management - Horses		4.40			0.49	0.08
4B8	Manure Management - Swine		26.70			17.85	0.53
4B9	Manure Management - Poultry		37.70			63.70	0.29
	Subtotal manure management		248.40	3.80		233.65	22.04
4D1	Direct Emissions from Agricultural Soils		35.60	81.31	29.34	190.81	26.07
4F	Field Burning of Agricultural Wastes		1.90		2.69	0.92	0.05
4G	Other Agricultural (including use of pesticides)			0.19			
5B	Forest and Grassland Conversion	0.20		0.01			
	Subtotal agricultural soils and wastes	0.20	1.90	0.20	2.69	0.92	0.05
6A	Solid Waste Disposal on Land		4.10				
6B	Waste Water Handling		5.50	3.92			5.03
6C	Waste Incineration	1.70		0.16			0.21
6D	Other Waste (incl. Composting and Biogas Prod.)	0.30				0.50	0.62
	Subtotal waste handling	2.00	9.60	4.08		0.50	5.87
7	Other		0.30				
z_5E	Other Sources and Sinks		8.30				
z_1A3ai(i)	International Aviation -Take-Off and Landing			0.08			
z_1A3ai(ii)	International Aviation - Cruise			1.04			
z_1A3di(i)	International Maritime and Inland Waterway Navigation			0.15			
	Total	1619.8	323.00	129.2	1281.5	442.1	125.9
					-20.9%	+36.9%	-2.6%

6.2.3 Observations for all regions

The overview comparisons of sectoral emission shares in the US and the PR of China (see Figs. 5 and 6.) indicate some substantial differences between the emission profiles of the US (and similarly Europe) on one side and China on the other. In particular the high contribution of emissions from agriculture to total NH₃ emissions in general is evident. In the case of NO_x emissions, mobile sources (road transport and off-road) contribute a larger share in Europe and the US, while power generation based on fossil fuels (coal mainly) in the energy

industries of China and the US contribute in a similar magnitude. A remarkable difference can be seen in the contribution of road transport to N_2O emissions and – to a smaller extent – NH_3 emissions in the US compared to Europe.

A comparison between the US NEI and EDGAR v4 indicates overall quite similar total figures, except for N_2O (EDGAR v4 substantially higher than US NEI). However, the sectoral structure for NO_x and NH_3 shows remarkable difference, for instance regarding the split between manure management and agricultural soils (NH_3) and power generation (NO_x).

6.3 Emission trends and implications

Apart from the analysis of emission inventories for specific years, which give a snapshot for a certain point in time, looking at the temporal trends in emissions can give valuable insight in the development of both emissions as well as the methodologies applied for their calculation.

Fig. 8 displays trends in NO_x emissions for the period 1990-2005, based on data from EMEP and the USEPA, and additional values from the literature for specific years or periods. A general downward trend can be observed for Europe and the US, reflecting significant emission control activities in particular with regard to stationary and mobile combustion sources. This trend occurs both in the official inventory figures (EMEP, USEPA) and the EDGAR data, with similar slopes, but with different starting points.




Fig. 8. Estimated trends in NH_3 (left) and NO_x (right) emissions in Europe, the US and the PR of China for the period 1990-2005

In the case of the People's Republic of China, no official dataset for a longer term trend analysis is yet available. Using a synthesis of satellite observations and bottom-up emission inventory development, Zhang et al. (2007) conclude that NO_x emissions in China from 1995-2004 likely increased by 70%. This is of particular relevance due to the amount of emissions, which – if emissions in China were set to further increase – may well exceed those of the US and Europe, especially while these show stagnating (NH_3) or downward trends (NO_x).

The picture for NH_3 emissions (Fig. 8) shows no significant reductions in the 15 year period indicated. A moderate downward trend in Europe can be observed (-18%), which is largely attributable to the political restructuring of Eastern Europe after 1989 (Horvath and Sutton, 1998) while US emissions have increased slightly in the period from 1990 to 2000. Although the values shown in Figure 6 indicate a small (6%) reduction below 1990 for 2005, this is due to a change in the calculation methodology in 2000, which has not been reapplied to the

earlier years. If the old methodology had been maintained throughout (with 2001 set equal to 2000), this would imply an overall national increase of about 22% between 1990 and 2005 comparing the US NEI values for both years. Again, Chinese emissions of NH_3 have as of yet only been estimated by few authors, with 1990 emission estimates ranging between 8 and 10 Tg, with a figure of 11.7 Tg (Klimont 2001b) for the year 1995. This latter figure would indicate an increase of 21% within these five years. This figure is attributed by Klimont (2001b) to a estimated 27% increase in NH_3 emissions from livestock and a 19% increase in fertilizer related emissions, reflecting increased consumption of livestock products in China, especially poultry (57% increase).

Emission trends for N_2O emissions are subject to significant uncertainties in the same way as the annual emission estimates. UNFCCC reported data for Europe and the US, as well as emission figures from the EDGAR Hyde 1.4 (Van Aardenne et al., 2001) inventory of historic emission trends and EDGAR v4 (EDGAR, 2008) emissions for the year 2000, which also include China make a rough trend assessment possible. For Europe, a consistent downward trend for both EDGAR Hyde 1.4 and UNFCCC figures can be observed, but it has to be noted that EDGAR FT32 (Olivier et al., 2005; Van Aardenne et al., 2005) for the year 2000 are significantly above UNFCCC figures, and higher than 1995 emissions from EDGAR Hyde 1.4. For the US, EDGAR Hyde 1.4 pictures a slow increase for the period 1990 to 1995, with 2000 emissions remaining at approximately the same level, while UNFCCC figures are slightly above $1.3 \text{ Tg N}_2\text{O yr}^{-1}$. A significant increase of N_2O emissions from the PR of China is portrayed in EDGAR Hyde 1.4, but EDGAR FT32 emissions for 2000 are lower than EDGAR Hyde for 1995, and not in the pathway that could be interpolated from the sharp increase of Chinese emissions in the period 1990 to 1995. Hence some caution is needed in considering the overall trends for the period 1990 to 2000.

7 Discussion and Conclusions

7.1 Completeness and coverage of emission inventories

The picture presented in the previous sections is quite diverse across different dimensions. In many cases, emission inventories clearly reflect the purpose they have been designed for, i.e. serving for regulatory purposes (e.g. the UNFCCC and UNECE CLRTAP EMEP inventories), where compliance with international protocols drives the need for a pragmatic accounting system for emissions. On the other hand, inventories such as the EDGAR emission database are bottom-up, science driven compilations of emissions based on emission factors and – typically – publicly available, statistical information on activity rates. While the legal implications and validation of national submissions are key aspects for the former, the latter have the main objective to provide comprehensive and consistent datasets for (atmospheric) modelling exercises. This is somewhat reflected as well in the sectoral structure in which these inventories are compiled.

In terms of completeness, two aspects need to be considered: firstly, are all known sources of emissions reflected in the inventory, or can missing sources be identified based on the state-of-the-art of scientific research? And secondly, have all known emissions been calculated based on a consistent methodology and no figures are missing or not estimated?

Officially reported emission inventory data are often incomplete with regard to both aspects, respectively are only available with a significant time lag due to the time it takes to compile them based on official statistics and to undergo validation and error checking, before they are officially submitted. Independent inventories however do not always have access to the latest, most accurate and detailed country specific EFs, penetration of measures and technology etc. for each country and hence are prone to over/underestimate emissions from individual sectors.

At the same time, there are clearly missing sources that have been identified and quantified in recent years, for instance emissions from NO from agricultural and forest soils and other natural and biogenic emissions (NH₃ from sea birds, NO_x from lightning and forest fires etc.). Compared to the main contributing sectors of anthropogenic emissions, the contributions of these missing sources to date is small, but with decreasing emissions due to emission control activities, their relative importance will continue to grow in the future. However, there may yet be major missing sources in the inventories. For example low temperature coal and other biomass combustion (Sutton et al., 2008) may be a major additional source of ammonia emissions in China that has yet to be quantified. Although of historical importance (Fowler et al. 2004), coal burning in residential combustion sources is no longer a significant activity for much of Europe and North America. Similarly, there remain major uncertainties in regional nitrous oxide emissions, as indicated by the assessment of Crutzen et al. (2007) who estimated from the global rate of N₂O increase that total agricultural emissions should be 4-5% of the input reactive nitrogen, which is larger than the base emission rate of 1% used in the IPCC methodology. This differential is most likely due to other biogenic sources of N₂O, but it cannot be ruled out that there are missing industrial or combustion source emissions.

In comparing the regulatory and science based inventories, it is possible to identify systematic errors or gaps in both, provided there is sufficient evidence and the level of detail provided allows for an in-depth assessment of the methodology used and factors and values applied.

The coverage of regions and substances for instance in the EDGAR inventory is comprehensive, but the fact that the NH₃ inventory for the time after 1990 is not yet available hints at the substantial resources and time required to compile these datasets. This is particularly problematic in the case of ammonia, as this is going to be a key pollutant to be targeted in future reduction efforts at least in Europe due to its growing relative importance

for the control of secondary aerosol concentrations and its contribution to acidification and eutrophication.

Finally, the spatial and temporal coverage of emission data as input to atmospheric dispersion models is a key issue. On a global scale, a resolution of 1×1 or 0.5×0.5 degree is sufficient to capture the general trends and concentrations/depositions. Yet, for regional and local modelling, inventories with a resolution of more than $10 \text{ km} \times 10 \text{ km}$ are likely to miss vital distribution patterns and thus lead to mismatches between model results and observations. And while national emission inventories with a spatial resolution of e.g. $1 \text{ km} \times 1 \text{ km}$ are available for some countries, they will rarely have been compiled using the same datasets and assumptions as the global inventories, hence may create boundary problems when nesting different inventories for modelling purposes (see as well Reis et al., 2008). The differences in the sectoral emissions splits between the inventories described in Section 6.2 have implications as well for the temporal and spatial profiles of these emissions when used as input for atmospheric dispersion models. Ultimately, this may lead to quite different ambient modelled concentrations of primary and secondary pollutants depending on which inventory data are used, in particular for pollutants such as tropospheric ozone, where the time and the location of the precursor emissions matters.

7.2 Trends and uncertainties

The trends for the three gases investigated in this paper show quite different patterns, as well as differences between the countries and regions included in this paper. NO_x emissions in Europe and the US have fallen markedly, reflecting successful emission control policies especially since the early 1990s. NO_x emissions from the PR of China, while much lower than those of Europe and the US, do not show a consistent trend due to the lack of data available. However, with the substantial growth of the Chinese economy and hence energy demand,

which is mainly supplied by coal fired power plants, a continuing increase of NO_x emissions can be anticipated.

Ammonia emission trends show a quite different picture, with only minor reductions in Europe over the period of 1990 to 2005, which were mainly a side effect of political restructuring around 1990, while the US ammonia emissions have increased by around 20% over the period. In both areas there have been major within-region differences. For example, in Europe, significant ammonia emission reductions were estimated for Denmark and the Netherlands, where control policies have been in place for more than a decade, while conversely emissions in Spain are estimated to have increased substantially over the last 20 years. In Europe there has been substantial debate as to whether estimated emission trends are verified by measurements (e.g. Horvath and Sutton, 1998; Sutton et al., 2003), and a recent assessment by Bleeker et al. (2008) concluded that the trend in reductions in ammonia emission in the Netherlands is supported by the measurements, though debate remains concerning the absolute magnitude of emissions. For the US, the overall increase in ammonia emissions is matched by major regional increases, especially in the Midwest (Sutton et al., 2008, based on Lehman et al., 2007) and in the Eastern US (Bleeker et al., 2008). With only a few data points available for China, it is yet evident that emissions are estimated to be about 2 to 3 times higher than European or US emissions in this period. In addition to that, Klimont (2001) projected a steep increase in NH₃ emissions for the period 1990-1995, equally matched between increased livestock numbers and mineral fertilizer use. Future projections for China suggest further increase, with Klimont (2001) estimating emissions in the range 8.2 to 19.9 Tg NH₃ for 2030.

Trends for nitrous oxide emissions finally indicate a slow decrease of emissions in Europe, a slight increase in US N₂O emissions, and a substantial growth in emissions in China. However, the trends depicted by data from the EDGAR Hyde 1.4 project and those reported

to UNFCCC do not match well with 2000 figures from EDGAR FT32, indicating different methodologies, respectively potentially different source sectors being included/excluded.

Some of the differences hint as well at the varying state of knowledge and certainty with regard to both the current amount and the future direction of emissions and trends. For European and US emissions, NO_x emission values can be seen as quite robust and well understood, while in particular the trend of NO_x emissions from China is highly uncertain due to the unknown developments in the Chinese energy demand and potential efficiency gains and decoupling efforts between economic growth and energy demand. This is to some extent also the case for European NO_x emissions. The phasing out of nuclear power on the one hand, and the plans to increase the use of domestic coal and biomass for power generation on the other hand make it more difficult to predict the development of NO_x emissions from the power generation sector beyond 2010 (Vestreng et al., 2008).

Ammonia emissions are dominated by agricultural production, and here future trends will be heavily influenced by agricultural policies, such as for instance the Common Agricultural Policy (CAP) Reform in Europe. Livestock numbers and the implementation of measures under the Integrated Pollution Prevention and Control (IPPC) directives will introduce further stringent measures to reduce NH_3 emissions from animal housing. More evidence is yet required to assess if China's ammonia emissions are indeed growing at a rate as the few data points suggest, for example illustrated in the wide range between the 2030 projections of Klimont (2001), the high estimate of which reflected a simple projection of activity statistics, while the low projection assumed full implementation of Best Available Techniques.

For N_2O , both the general uncertainty about the emissions, and their trends, are significantly larger than for the other pollutants. How trends after 2010 evolve will depend on future negotiations and the way how Kyoto and post-Kyoto greenhouse gas emission targets are achieved, and how further reductions will be set.

7.3 Future research needs

A first step to reduce the uncertainties identified above and to support the efforts to close the gaps in N budgets on a global or regional scale will be to detect missing sources and to quantify their contribution to overall emissions. This is of particular importance in the field of natural and semi-natural sources of emissions, which are often not yet well covered by emission inventories, which focus on anthropogenic emissions due to their history and heritage. Current research targeting unknown emission sources on the one hand, and recent achievements in quantifying emissions e.g. of NO from agricultural and forest soils (Skiba et al., 2001; Friedrich et al., 2007), NH₃ from seabirds and other animals (Blackall et al., 2007) need to find their way into the guidelines for emission reporting and become a regular part of emission inventories. At the same time further efforts are needed to better quantify important but highly uncertain emissions such as ammonia from combustion sources (Fowler et al., 2004; Sutton et al., 2008) and apparently missing sources of nitrous oxide (Crutzen et al., 2007). Only if inventories reflect the current state-of-the-art of research can validation and verification experiments, comparing model calculations with observations on different spatial scales (e.g. Bleeker et al., 2008; Bergamaschi et al. 2005; Henze et al., 2008) deliver meaningful results towards a better understanding of the processes through which reactive and non-reactive nitrogen species contribute to air pollution and climate change. In addition, a fully quantified nitrogen cycle is a key requirement for the design of intelligent and effective management options towards reducing the negative impacts of N along the full nitrogen cascade. Complete and accurate emission inventories of all nitrogen species are a crucial building block for this.

8 Acknowledgements

Most of the work described here has been supported by a range of research projects and ongoing activities, such as the European Commission funded research project NitroEurope IP,

COST Action 729 the ESF Programme Nitrogen in Europe (NinE). The present paper is a contribution to activities under the International Nitrogen Initiative (INI) and the National Basic Research Program of China (Grant No. 2007CB407303). The US EPA through its Office of Research and Development (partially funded and collaborated in) the research described here. It has been subjected to Agency review and approved for publication.

9 References

- Akimoto, H., and Narita, H.: Distribution of SO₂, NO_x and CO₂ emission from fuel combustion and industrial activities in Asia with 1°×1° resolution, *Atmospheric Environment*, 28, 213-225, 1994.
- Bai, N. B.: The emission inventory of SO₂, NO_x and CO₂ in China. The Atmospheric Ozone Variation and Its Effect on the Climate and Environment in China, China Meteorological Press, 145-150, 1996.
- Bergamaschi, P., Krol, M., Dentener, F., Vermeulen, A., Meinhardt, F., Graul, R., Ramonet, M., Peters, W., and Dlugokencky, E. J.: Inverse modelling of national and European CH₄ emissions using the atmospheric zoom model TM5. *Atmos. Chem. Phys.*, 5:2431-2460, 2005
- Bishop, G. A., and Stedman, D. H.: A Decade of On-road Emission Measurements, *Environmental Science and Technology*, 42, 1651-1656, 2008.
- Blackall, T. D., Wilson, L. J., Theobald, M. R., Milford, C., Nemitz, E., Bull, J., Bacon, P. J., Hamer, K. C., Wanless, S., and Sutton, M. A.: Ammonia emissions from seabird colonies, *Geophysical Research Letters*, 34, L10801, doi:10.1029/2006GL028928, 2007.
- Bleeker, A., Sutton, M. A., Acherman, B., Alebic-Juretic, A., Aneja, V. P., Ellermann, T., Erismann, J. W., Fowler, D., Fagerli, H., Gauger, T., Harlen, K. S., Hole, L. R., Horváth, L., Mitosinkova, M., Smith, R. I., Tang, Y. S., and van Pul, A.: Linking ammonia emission trends to measured concentrations and deposition of reduced nitrogen at different scales, in: *Atmospheric Ammonia: Detecting emission changes and environmental impacts*, edited by: M.A. Sutton, S. R. a. S. B., Springer, New York, 2008.
- Bouwman, A. F., Lee, D. S., Asman, W. A. H., Dentener, F. J., Van der Hoek, K. W., and Olivier, J. G. J.: A global high-resolution emission inventory for ammonia, *Global Biogeochemical Cycles*, 11, 561-587, 1997.
- Bradshaw, J., Davis, D., Grodzinsky, G., Smyth, S., Newell, R., Sandholm, S., and Liu, S.: Observed distribution of nitrogen oxides in the remote free troposphere from the NASA global tropospheric experiment programs, *Reviews of Geophysics* 38, 111-116, 2000.
- Brasseur, G. P., Orlando, J. J., and Tyndall, G. S.: *Atmospheric Chemistry and Global Change*, Oxford University Press, New York, 1999.
- Crutzen, P. J., Mosier, A. R., Smith, K. A., and Winiwarter, W.: N₂O release from agro-biofuel production negates global warming reduction by replacing fossil fuels, *Atmos. Chem. Phys. Discuss.*, 7, 11191-11205, 2007.
- Del Grosso, S. J., Parton, W. J., Mosier, A. R., Hartman, M. D., Brenner, J., Ojima, D. S., Schimel, D. S., "Simulated Interaction of Carbon Dynamics and Nitrogen Trace Gas Fluxes Using the DAYCENT Model." In Schaffer, M., Ma, L., Hansen, S. (eds.); *Modeling Carbon and Nitrogen Dynamics for Soil Management*. CRC Press. Boca Raton, FL. 303-332. 2001.
- Dragosits, U.; Dore, A.; Sheppard, L.; Vieno, M.; Tang, Y. S.; Theobald, M.; Sutton, M.: Sources, dispersion and fate of atmospheric ammonia. In: Hatfield, J.L.; Follett, R.F., (eds.) *Nitrogen in the Environment - Sources, Problems, and Management*. Academic Press, 333-393, 61pp, 2008
- EDGAR: Emissions Database for Global Atmospheric Research (EDGAR): <http://edgar.jrc.ec.europa.eu/>, access: 22/12/2008, 2008.
- EMEP Webdab emission data hosted by the Centre on Emission Inventories and Projections (CEIP): <http://www.ceip.at/>, access: 22/12/2008, 2008.
- Erismann, J.W.; Mosquera, J., and Hensen, A.: Two options to explain the ammonia gap in The Netherlands, *Environmental Science & Policy*, 4 (2-3), 97-105, 2001.
- Erismann, J. W.: The Nanjing Declaration on Management of Reactive Nitrogen, *BioScience*, 54, 286-287, 2004.
- Erismann, J. W., Spranger, T., Sutton, M. A., Akselsson, C., Amin-Hanjani, S., Anderson, H. V., ApSimon, H., Fagerli, H., Harmens, H., Havlikova, M., Hettelingh, J. P., Hicks, K., Horvath, L., Hutchings, N. J., Maasikimets, M., Maione, M., Reis, S., and Stenby, C.: Working Group 5 : Nitrogen – integrated environmental policies. *Proceedings of the UNECE/EC Saltsjobaden 3 Workshop*, 2007.
- Erismann, J. W., Sutton, M. A., Galloway, J. N., Klimont, Z., and Winiwarter, W.: 100 years of ammonia synthesis: how a single patent changed the world. , *Nature Geoscience*, 1, 636 - 639, 2008.

- Fowler, D., O'Donoghue, M., Muller, J. B. A., Smith, R. I., Dragosits, U., Skiba, U., Sutton, M. A., and Brimblecombe, P.: A Chronology of Nitrogen Deposition in the UK between 1900 and 2000, *Water, Air and Soil Pollution: Focus*, 4, 9-23, 2004.
- Friedrich, R., and Theloke, J.: Improving and applying methods for the calculation of natural and biogenic emissions and assessment of impacts to the air quality 2007.
- Galloway, J. N.: The global nitrogen cycle: changes and consequences, *Environmental Pollution*, 102, 15-24, 1998.
- Galloway, J. N., Aber, J. D., Erisman, J. W., Seitzinger, S. P., W., H. R., Cowling, E. B., and Cosby, B. J.: The nitrogen cascade, *BioScience*, 53, 341-356, 2003.
- Galloway, J. N., Dentener, F. J., Capone, D. G., Boyer, E. W., Howarth, R. W., Seitzinger, S. P., Asner, G. P., Cleveland, C. C., Green, P. A., Holland, E. A., Karl, D. M., Michaels, A. F., Porter, J. H., Townsend, A. R., and Vorosmarty, C. J.: Nitrogen cycles: past, present and future, *Biogeochemistry*, 70, 153-226, 2004.
- Galloway, J. N., Townsend, A. R., Erisman, J. W., Bekunda, M., Cai, Z., Freney, J. R., Martinelli, L. A., Seitzinger, S. P., and Sutton, M. A.: Transformation of the Nitrogen Cycle: Recent Trends, Questions and Potential Solutions, *Science*, 320, 889-892, 2008.
- Gilliland, A. B., Appel, K. W., Pinder, R. W., and Dennis, R. L.: Seasonal NH₃ Emissions for the Continental United States: Inverse Model Estimation and Evaluation, *Atmospheric Environment*, 40 4986-4998, 2006.
- Godowitch, J. M., Hogrefe, C., and Rao, S. T.: Diagnostic Analyses of a Regional Air Quality Model: Changes in Modeled Processes Affecting Ozone and Chemical-Transport Indicators from NO_x Point Source Emission Reductions, *J. Geophysical Research*, in press, 2008.
- Goebes, M. D., Strader, R., Davidson, C. An ammonia emission inventory for fertilizer application in the United States, *Atmospheric Environment*, 37, 2539-2550, 2003.
- Henze, D. K., Seinfeld, J. H., and Shindell, D. T.: Inverse modeling and mapping US air quality influences of inorganic PM_{2.5} precursor emissions using the adjoint of GEOS-Chem, *Atmospheric Chemistry and Physics Discussions*, 8, 15031-15099, 2008.
- Hettelingh, J. P., Posch, M., Slootweg, J., and (eds.): CCE Status Report 2008, Coordination Centre for Effects (CCE), 2008.
- Horváth, L., and Sutton, M. A.: Long term record of ammonia and ammonium concentrations at K-puszt, Hungary, *Atmospheric Environment*, 32, 339-344, 1998.
- IPCC: Guidelines for National Greenhouse Gas Inventories: Greenhouse Gas Inventory Reference Manual (Revised 1996), IPCC/OECD/IGES, Bracknell, UK, 1997.
- IPCC: Good Practice Guidance, Uncertainty Management in National Greenhouse Gas Inventories, IPCC/IGES, Kanagawa, Japan, 2000.
- Kato, N., and Akimoto, H.: Anthropogenic emissions of SO₂ and NO_x in Asia: Emission inventories, *Atmospheric Environment*, 26A, 2997-3017, 1992.
- Kim, S. W., Heckel, A., McKeen, S. A., Frost, G. J., Hsie, E. Y., Trainer, M. K., Richter, A., Burrows, J. P., Peckham, S. E., and Grell, G. A.: Satellite-observed US power plant NO_x emission reductions and their impact on air quality, *Geophysical Research Letters*, 33, L22812, 2006.
- Kim, S.-W., Heckel, A., Frost, G. J., Richter, A., Gleason, J., Burrows, J. P., McKeen, S., Hsie, E.-Y., Granier, C. and Trainer M. NO₂ columns in the western United States observed from space and simulated by a regional chemistry model and their implications for NO_x emissions, *J. Geophys. Res.*, 114, D11301, doi:10.1029/2008JD011343, 2009.
- Klaassen, G.: Options and Costs of Controlling Ammonia Emissions in Europe, *Agricultural Economics*, 21, 219-240, 1994.
- Klimont, Z.: Current and Future Emissions of Ammonia in China. 10th International Emission Inventory Conference - "One Atmosphere, One Inventory, Many Challenges", US EPA, <http://www.epa.gov/ttn/chief/conference/ei10/ammonia/klimont.pdf>, access: 22/12/2008, 2001a.
- Klimont, Z.: Ammonia Emissions, Abatement Technologies and Related Costs for Europe in the RAINS model, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria, 2001b.

- Lee, D. S., Köhler, I., Grobler, E., Rohrer, F., Sausen, R., Gallardo-Klenner, L., Olivier, J. G. J., Dentener, F. J., and Bouwman, A. F.: Estimations of global NO_x emissions and their uncertainties, *Atmospheric Environment*, 31, 1735–1749, 1997.
- C.M.B. Lehmann, V.C. Bowersox, R.S. Larson and S.M. Larson: Monitoring long-term trends in sulphate and ammonium in US precipitation: results from the National Atmospheric Deposition Program/National Trends Network, *Water Air Soil Pollut. Focus* 7, 1567–7230, 2007; (http://bqs.usgs.gov/acidrain/Deposition_trends.pdf) (accessed 1 Nov. 2007).
- Li, Y. E., and Lin, E.: Emissions of N_2O , NH_3 and NO_x from fuel combustion, industrial processes and the agricultural sectors in China, *Nutrient Cycling in Agroecosystems*, 57, 99–106, 2000.
- Li, C., Zhuang, Y., and Cao, M.: Comparing a national inventory of N_2O emissions from arable lands in China developed with a process-based agroecosystem model to the IPCC methodology, *Nutrient Cycling in Agroecosystems*, 60, 159–170, 2001.
- Lu, Y. Y., Huang, Y., Zou, J. W., and Zheng, X. H.: An inventory of N_2O emissions from agriculture in China using precipitation-rectified emission factor and background emission., *Chemosphere* 65, 1915–1924, 2006.
- Ma, J., and Zhou, X. J.: Development of a three dimensional inventory of aircraft NO_x emissions over China. , *Atmospheric Environment*, 34, 389–396, 2000.
- Mosier, A. R., Duxbury, J. M., Freney, J. R., Heinemeyer, O., and Minami, K.: Assessing and mitigating N_2O emissions from agricultural soils, *Climatic Change*, 40, 7–38, 1998.
- NARSTO: Improving Emission Inventories for Effective Air Quality Management across North America, 2005.
- Olivier, J. G. J., Bouwman, A. F., Berdowski, J. J. M., Veldt, C., Bloos, J. P. J., Visschedijk, A. J. H., Van der Maas, C. W. M., and Zandveld, P. Y. J.: Sectoral emission inventories of greenhouse gases for 1990 on a per country basis as well as on $1^\circ \times 1^\circ$, *Environmental Science & Policy*, 2, 241–264, 1999.
- Olivier, J.G.J. and J.J.M. Berdowski Global emissions sources and sinks. In: Berdowski, J., Guicherit, R. and B.J. Heij (eds). "The Climate System", 33-78. A.A. Balkema Publishers/Swets & Zeitlinger Publishers, Lisse, The Netherlands, 2001.
- Olivier, J.G.J., Van Aardenne, J.A., Dentener, F., Ganzeveld, L. and J.A.H.W. Peters.: Recent trends in global greenhouse gas emissions: regional trends and spatial distribution of key sources. In: "Non- CO_2 Greenhouse Gases (NCGG-4)", A. van Amstel (coord.), page 325-330. Millpress, Rotterdam, ISBN 90 5966 043 9, 2005
- Parrish, D. D.: Critical evaluation of US on-road vehicle emission inventories, *Atmospheric Environment*, 40, 2288-2300, 2006.
- Pinder, R. W., Adams, P. J., Pandis, S. N., and Gilliland, A. B.: Temporally Resolved Ammonia Emission Inventories: Current Estimates, Evaluation Tools, and Measurement Needs, *J. Geophysical Research*, 111 D16310, 2006.
- Press, A.: China Agricultural Yearbook (in Chinese), Agricultural Press, Beijing, 1996.
- Reis S., Pfeiffer H., Scholz Y. and Theloke J.: Temporal and Spatial Distribution of Carbon Emissions. In: Dolman, H (ed) The Continental-Scale Greenhouse Gas Balance of Europe. Springer Publishers, 2008
- Richter, A., Burrows, J.P., Nüß, H., Granier, C., and Niemeier, U.: Increase in tropospheric nitrogen dioxide levels over China observed from space, *Nature*, 437, 129-132, 2005
- Rypdal, K., and Winiwarter, W.: Uncertainties in Greenhouse Gas Inventories - Evaluation, comparability and implications. , *Environmental Science and Policy*, 4, 107–116, 2001.
- Skiba, U., Sozanska, M., Metcalfe, S., and Fowler, D.: Spatially disaggregated inventories of soil NO and N_2O emissions for Great Britain., *Water Air & Soil Pollution*, 1, 2001.
- Streets, D. G., and Waldhoff, S. T.: Present and future emissions of air pollutants in China: SO_2 , NO_x and CO , *Atmospheric Environment*, 34, 2000.
- Streets, D. G., Bond, T. C., Carmichael, G. R., Fernandes, S. D., Fu, Q., He, D., Klimont, Z., Nelson, S. M., Tsai, N. Y., Wang, M. Q., Woo, J.-H., and Yarber, K. F.: An inventory of gaseous and primary aerosol emissions in Asia in the year 2000, *Journal of Geophysical Research*, 108, 8809, 2003.

- Sutton, M.A., Dragosits, U., Tang, Y.S., and Fowler, D.: Ammonia emissions from non agricultural sources in the UK. *Atmos. Environ.* 34 (6), 855-869, 2000.
- Sutton, M. A., Asman, W. A. H., Ellerman, T., van Jaarsveld, J. A., Acker, K., Aneja, V., Duyzer, J. H., Horvath, L., Paramonov, S., Mitosinkova, M., Tang, Y. S., Achermann, B., Gauger, T., Bartnicki, J., Nefel, A., and Erisman, J. W.: Establishing the link between ammonia emission control and measurements of reduced nitrogen concentrations and deposition, *Environmental Monitoring and Assessment*, 82, 149-185, 2003.
- Sutton, M. A., Nemitz, E., Erisman, J. W., Beier, C., Butterbach Bahl, K., Cellier, P., de Vries, W., Cotrufo, M. F., Skiba, U., and al., c.: Challenges in quantifying biosphere-atmosphere exchange of nitrogen species, *Environmental Pollution*, 150, 125-139, 2007.
- Sutton, M. A., Erisman, J. W., Dentener, F., and Moeller, D.: Ammonia in the environment: from ancient times to the present, *Environmental Pollution*, 154, 2008.
- Tian, H. Z., Hao, J. M., Lu, Y. Q., and Zhu, T. L.: The NO_x emission distribution and inventory in China, *China Environmental Science*, 21, 493-497, 2001.
- UNEP: The Nanjing Declaration on Nitrogen Management Presented to the United Nations Environment Programme, 16th of October 2004, UNEP, Nanjing, PR of China, 2004.
- UNFCCC: Greenhouse Gas Inventory Data: http://unfccc.int/ghg_data/items/3800.php, access: 22/12/2008, 2008.
- USDA: Agricultural Projections to 2016, USDA 2007.
- USEPA: National Emission Inventory Tier Summaries: <http://www.epa.gov/ttn/chief/eiinformation.html>, access 14/08/2009, 2009a
- USEPA: Inventory of US Greenhouse Gas Emissions and Sinks: 1990-2007, <http://epa.gov/climatechange/emissions/usinventoryreport.html> access: 14/08/2009, 2009b.
- Van Aardenne, J. A., Carmichael, G. R., Levy II, H., Streets, D., and Hordijk, L.: Anthropogenic NO_x emissions in Asia in the period 1990–2020, *Atmospheric Environment*, 33, 633–646, 1999.
- Van Aardenne, J.A., Dentener, F.J., Olivier, J.G.J., Klein Goldewijk, C.G.M. and J. Lelieveld: A 1 x 1 degree resolution dataset of historical anthropogenic trace gas emissions for the period 1890-1990. *Global Biogeochemical Cycles*, 15(4), 909-928, 2001.
- Van Aardenne, J.A., F. Dentener, J.G.J. Olivier and J.A.H.W. Peters: The EDGAR 3.2 Fast Track 2000 dataset (32FT2000), 2005
- van der A, R.J., Peters, D.H.M.U., Eskes, H., Boersma, K.F., Van Roozendael, M., De Smedt, I., and Kelder, H.M.: Detection of the trend and seasonal variation in tropospheric NO₂ over China, *J. Geophys. Res.*, 111, D12317, doi:10.1029/2005JD006594, 2006.
- Velthof, G. L., Oudendag, D., Witzke, H. P., Asman, W. A. H., Klimont, Z., and Oenema, O.: Integrated assessment of nitrogen emissions from agriculture in EU-27 using MITERRA EUROPE, *Journal of Environmental Quality*, submitted, 2008.
- Vestreng, V., Ntziachristos, L., Semb, A., Reis, S., Isaksen, I. S. A., and Tarrasón, L.: Evolution of NO_x emissions in Europe with focus on road transport control measures, *Atmos. Chem. Phys. Discuss.*, 8, 10697-10747, 2008.
- Wallace, J. M., and Hobbes, P. V.: *Atmospheric Science – An Introductory Survey*. 2nd Edition, Elsevier Publishers, 2006.
- Winiwarter, W., and Rypdal, K.: Assessing the uncertainty associated with national greenhouse gas emission inventories: a case study for Austria, *Atmospheric Environment*, 35, 5425–5440, 2001.
- Xing, G. X.: N₂O emission from cropland in China, *Nutrient Cycling in Agroecosystems* 52, 249–254, 1998.
- Yan, X. Y., Akimoto, H., and Ohara, T.: Estimation of nitrous oxide, nitric oxide and ammonia emissions from croplands in East, Southeast and South Asia, *Global Change Biology*, 9, 1080–1096, 2003.
- Zhao, D., and Wang, A.: Estimation of Anthropogenic Ammonia Emissions in Asia, *Atmospheric Environment*, 28, 689–694, 1994.

- Zheng, X. H., Han, S. H., Huang, Y., Wang, Y. S., and Wang, M. X.: Re-quantifying the emission factors based on field measurements and estimating the direct N₂O emission from Chinese croplands, *Global Biogeochemical Cycles*, 18, doi:10.1029/2003GB002167, 2004.
- Zhang, Q., Streets, D. G., He, K., Wang, Y., Richter, A., Burrows, J. P., Uno, I., Jang, C. J., Chen, D., Yao, Z., Lei, Y.: NO_x emission trends for China, 1995-2004: The view from the ground and the view from space, *J. Geophys. Res.*, 112, D22306, doi:10.1029/2007JD008684, 2007