

PON1 status of farmworker mothers and children as a predictor of organophosphate sensitivity

Clement E. Furlong^{a,*}, Nina Holland^{b,*}, Rebecca J. Richter^a, Asa Bradman^b, Alan Ho^b and Brenda Eskenazi^b

The objective was to determine PON1 status as a predictor for organophosphorus insecticide sensitivity in a cohort of Latina mothers and newborns from the Salinas Valley, California, an area with high levels of organophosphorus insecticide use. PON1 status was established for 130 pregnant Latina women and their newborns using a high-throughput two substrate activity/analysis method which plots rates of diazoxon (DZO) hydrolysis against rates of paraoxon (PO) hydrolysis. Arylesterase activity (AREase) was determined using phenylacetate as a substrate, allowing comparison of PON1 levels across *PON1*₁₉₂ genotypes in mothers and children. Phenylacetate hydrolysis is not affected by the Q192R polymorphism. Among newborns, levels of PON1 (AREase) varied by 26-fold (4.3–110.7 U/ml) and among mothers by 14-fold (19.8–281.4 U/ml). On average, children's PON1 levels were four-fold lower than the mothers' PON1 levels ($P < 0.001$). Average PON1 levels in newborns were comparable with reported hPON1 levels in transgenic mice expressing human PON1_{Q192} or PON1_{R192}, allowing for prediction of relative sensitivity to chlorpyrifos oxon (CPO) and DZO. The predicted range of variability in sensitivity of mothers and children in the same Latino cohort was 65-fold for DZO and 131 to 164-fold for CPO. Overall, these findings indicate that many of the newborns and some of

the mothers in this cohort would be more susceptible to the adverse effects of specific organophosphorus pesticide exposure due to their PON1 status. Of particular concern are exposures of pregnant mothers and newborns with low PON1 status. *Pharmacogenetics and Genomics* 16:183–190 © 2006 Lippincott Williams & Wilkins.

Pharmacogenetics and Genomics 2006, 16:183–190

Keywords: children, chlorpyrifos, diazinon, Latino cohort, paraoxonase 1 (PON1), PON1 status, pregnancy

^aDepartments of Genome Sciences and Medicine, Division of Medical Genetics, University of Washington, Seattle, Washington and ^bCenter for Children's Environmental Health, School of Public Health, University of California, Berkeley, California, USA

Correspondence and requests for reprints to Clement Furlong, University of Washington, Division of Medical Genetics, Box 357720, Seattle, WA 98195-7720, USA
Tel: +1 206 543 1193; fax: +1 206 543 3050;
e-mail: clem@u.washington.edu

Sponsorship: This study was supported by 2 P01 ES009605, ES11387, ES09883, P30 ES01896, EPA-R82670901-5, P60 MD00222, and NIEHS ES09601/EPA: RD-83170901. The contents of this paper are solely the responsibility of the authors and do not necessarily represent official views of the NIH, or the EPA.

Received 26 July 2005 Accepted 13 October 2005

Introduction

Recent biological and ambient monitoring data have indicated widespread organophosphate pesticide exposures to the US population, including adults, pregnant women, children and fetuses [1–9]. In some cases, these exposures may exceed health-based reference levels [10–11]. Although many uses, including residential applications, of chlorpyrifos (CPS) and diazinon (DZ) were recently restricted by agreements with registrants [12,13], agricultural uses are still widespread. Organophosphate exposures at high doses have profound effects, primarily on the central nervous system [14], and there is growing evidence in animals and humans to suggest that chronic low level exposure may affect neurodevelopment [5,15–17].

The young of many species are more susceptible to organophosphate toxicity than adults [18–22]. For example, the maximum tolerated dose (MTD) of CPS in

7-day-old (PND7) rats is approximately 7.7% of the MTD in adult animals [22]. One factor contributing to the increased sensitivity in newborns is that levels of paraoxonase 1/arylesterase (PON1), a key organophosphate detoxifying enzyme, are three- to four-fold lower than in adults [23–27]. Even among adults, the levels of PON1 can vary by at least 13-fold [28].

Human PON1 enzyme, a high-density lipoprotein-(HDL) associated esterase, is encoded as a 355 amino acid protein by the *PON1* gene on chromosome 7q21.3–22.1 [29] with only its initiator methionine residue removed before incorporation into HDL particles [30,31]. In humans, a Q192R polymorphism affects the catalytic efficiency of hydrolysis of some organophosphate substrates [28,32,33], including chlorpyrifos oxon (CPO), the toxic metabolite of CPS [34]. The characterization of all 28 TagSNPs accounts for only 28% of the variance in PON1 levels (G.P. Jarvik, personal communication), much of which is attributable to a C-108T polymorphism in an Sp1 binding site in the 5' regulatory region [35–37].

*The first two authors contributed equally to this work.

Simple biochemical principles dictate that rates of detoxication of substrates are dependent on enzyme levels. Thus, when considering the effects of genetic variability on sensitivity to CPS/CPO exposures, both the quantity (level) of PON1 as well as the quality (Q192 versus R192) of PON1 must be considered. In other cases where the catalytic efficiency of hydrolysis of the two PON1₁₉₂ alloforms is equivalent [e.g. for diazoxon (DZO) hydrolysis], it is the level of PON1 that is important [34].

Li *et al.* [38] introduced the term PON1 status to include both PON1 level and functional *PON1*₁₉₂ genotype (Q/Q; Q/R; R/R) [38]. In this method, PON1 status is determined with a simple, high-throughput two-substrate assay and analysis where rates of DZO hydrolysis are plotted against rates of PO hydrolysis using serum or non-EDTA preserved plasma samples. This method, validated in adults, provides a functional determination of the PON1₁₉₂ alloform(s) present in plasma, as well as the level of an individual's plasma PON1 [39–41], both important in modulating exposures to organophosphorus compounds as well as other risks associated with PON1 status. The lower PON1 levels associated with PON1_{M55} [42] are primarily attributable to linkage disequilibrium with the inefficient promoter polymorphism *T-108* [35].

The development of animal model systems has provided important insights into the role of PON1 in detoxifying specific organophosphorus compounds. Injection of purified rabbit PON1 into rats increased resistance to paraoxon (PO) exposure [43,44] and, more significantly, to CPO exposure [44] because rabbit PON1 hydrolyses CPO very rapidly [45]. These observations were confirmed and extended using mice, which required much less purified PON1 for injections and were more amenable to genetic manipulation [34,38,46]. Injected PON1 protected against CPS/CPO exposures when injected 30 min or 24 h before exposure or up to 3 h post exposure [47]. These experiments provided convincing evidence that high levels of PON1 protected against CPS/CPO exposures. The increase in resistance was most dramatic against CPO exposures [34].

The generation of *PON1* knockout mice by Shih *et al.* [46] provided a model with which to examine the consequence of the absence of plasma PON1 on resistance to organophosphate exposures. The PON1 null mice were found to have dramatically increased sensitivity to CPO [46] and DZO exposure [34], and less noticeably increased sensitivity to the respective parent compounds CPS [46,47] and DZ [34]. It was surprising to find that the PON1 null mice did not have increased sensitivity to PO exposure [34].

Injection of purified human PON1_{Q192} or PON1_{R192} reconstituted PON1 activity in the serum of the *PON1*

null mice allowed for the testing of the efficiency of each human PON1₁₉₂ alloform in protecting against exposure under physiological conditions without exposing human subjects to these toxic organophosphates [34]. Examination of the catalytic efficiencies of hydrolysis of CPO, DZO and PO showed that it was the catalytic efficiency that determined whether PON1 would protect against exposures. Either hPON1₁₉₂ alloform protected equally well against DZO exposure, in agreement with the equivalent catalytic efficiency of each alloform for DZO hydrolysis. However, the hPON1_{R192} alloform provided significantly better protection against CPO exposure, in agreement with the higher catalytic efficiency of PON1_{R192} for CPO hydrolysis [34]. These observations on CPO exposures have been confirmed in transgenic mice expressing one or the other PON1₁₉₂ alloform at equivalent levels [48]. The finding that PON1_{Q192} does not protect as well as PON1_{R192} against CPO/CPS exposures is important because up to 50% of the general population is homozygous for PON1_{Q192} [49]. The catalytic efficiency for PO hydrolysis was too low to provide protection against PO exposures [34].

In the present study, we determined the PON1 status of 130 pregnant Latina women and their newborns living in an agricultural community in California [7], a region where approximately 22 727 kg of organophosphates are used annually [50]. We have previously reported that maternal urinary dialkyl phosphate metabolite levels are higher in this population relative to national reference data [8] and were associated with a shorter gestational age [51] and an increased frequency of abnormal reflexes in neonates [16]. The main aims of this study were to assess PON1 status in newborns and mothers and to predict their relative sensitivity to specific organophosphorus insecticides based on recent studies with 'PON1 humanized transgenic mice' expressing either human PON1_{R192} (hPON1₁₉₂) or PON1_{Q192} (hPON1_{Q192}) at equivalent levels [48].

Methods

Subjects and recruitment

A subset of 130 maternal-newborn pairs were randomly selected from the CHAMACOS cohort (Center for the Health Assessment of Mothers and Children of Salinas), a longitudinal birth cohort study ($n = 601$ enrollees, 528 live births) of the effects of environmental exposures on the health of children living in the Salinas Valley [7]. Women were eligible for enrollment in the CHAMACOS study if they were 18 years or older, less than 20 weeks gestation at enrollment, English- or Spanish-speaking, Medi-Cal eligible, and planning to deliver at the Natividad Medical Center. All women were Latina by ethnicity, including 87% born in Mexico, and the remainder in the USA. Approximately 28% of the women had worked in the fields during the pregnancy and

another 14% had other jobs in agriculture, including packing shed, nursery and greenhouse work. Overall, 82% of subjects had agricultural workers living in their homes during pregnancy. Additional information about exposure and associations with health outcomes is reported elsewhere [7,8,16,51]. All study protocols were approved by both the University of California, Berkeley and the University of Washington human-subject review processes. Informed consent was obtained for all subjects.

Biological sample collection and processing

Blood was collected from mothers at the time of their glucose tolerance test (26 ± 2.3 weeks' gestation) and in the hospital immediately before or after delivery. Umbilical cord blood was collected by delivery room staff once the baby was safely delivered. Heparinized whole blood was centrifuged, and divided into plasma, buffy coats and red blood cells, and stored at -80°C . Processed plasma samples were stored at -80°C before being shipped on dry ice to the University of Washington, Seattle, where they were also stored at -80°C until analysis of enzyme activity. We first compared PON1 activities in 25 mothers at two time points (26 weeks gestation and delivery). PON1 activities were highly correlated between the two time points for all three measured activities of PON1 ($r = 0.7, 0.8, 1.0, P < 0.0001$ for AREase, CPOase and DZOase, respectively). The ranges for the two measurements were comparable. Therefore, in subsequent samples, we measured PON1 activities in blood collected at one time point only (26 weeks' gestation).

Determination of PON1 status

PON1 status (functional *PON1*₁₉₂ genotype and plasma level) was determined by the method developed and validated on adult populations by Richter and Furlong [39]. This method provides an accurate determination of functional *PON1*₁₉₂ genotype through the use of a two-substrate enzyme kinetic analysis. PON1 enzyme activities in plasma of mothers and children were measured with three different substrates, including paraoxon (POase), diazoxon (DZOase) and phenylacetate (AREase), according to published protocols [39–41,52]. Because phenylacetate hydrolysis is not affected by the Q192R polymorphism and has been shown to correspond with PON1 levels determined by immunological methods [29,53], AREase activity was used for measurement of PON1 levels across genotypes. For PON1 status determination, rates of DZO hydrolysis were plotted against rates of PO hydrolysis for each mother and newborn child (cord blood) in the study. This analysis separates the population into three distinct groups, individuals functionally homozygous for PON1_{Q192}, PON1_{Q/R192} heterozygotes and individuals functionally homozygous for PON1_{R192}. The accuracy of the *PON1*₁₉₂ functional genotype determination has been verified by polymerase chain reaction (PCR) analysis of more than 2000 adult

samples and has been shown to identify individuals with mutations in the PON1 gene [41]. Although this plot suggests that PON1_{R192} homozygotes have lower rates of DZO hydrolysis than heterozygotes and PON1_{Q192} homozygotes, this is not the case, as the position 192 alloforms have equivalent catalytic efficiencies for DZO hydrolysis [34]. However, the PON1_{R192} alloform is more sensitive to inhibition by the high salt concentration intentionally used in this assay to resolve the three PON1₁₉₂ phenotypes. The distributions and descriptive statistics of PON1 status and enzyme levels in mothers and their newborns were analysed by STATA 8.0 [54].

Results


PON1 status

Figure 1 shows the PON1 status for mothers and newborns (cord blood) as determined by the plot of rates of hydrolysis of DZO versus PO. This plot clearly resolves the three functional PON1₁₉₂ phenotypes (verified by PCR analysis) for the mothers. However, a number of data points for newborns with lower activity values (22 out of 130 or approximately 17%) were not resolved by this analysis and could only be accurately assigned by genotyping the Q192R polymorphism. However, the PON1 status plot provided relative PON1 levels for each of the 22 subjects. This analysis also shows the effects of independent allelic expression in the heterozygotes (including the newborns) with some individuals expressing more of one PON1₁₉₂ alloform than the other. Genotyping alone does not provide this information. Several points are worth noting from this analysis: (i) there is a large variability of PON1 activity values within each *PON1*₁₉₂ genotype, both for mothers and newborns; (ii) both groups of homozygotes fall closely to the trend line; and (iii) the heterozygote values vary considerably from the trend line (trend line not shown), consistent with the concept of independent cis regulation of each *PON1* allele (Fig. 1).

Range of plasma PON1 levels


Figure 2 shows PON1 levels across genotypes for mothers and newborns estimated by the AREase activity. Figure 3 shows the effects of the *PON1*₁₉₂ polymorphism on rates of paraoxon hydrolysis across genotypes for mothers and newborns, and provides an excellent example of why substrates whose rates of hydrolysis are affected by the *PON1*₁₉₂ polymorphism should not be used to compare levels across position 192 genotypes. Rates of paraoxon hydrolysis are significantly lower for Q/Q homozygotes and heterozygotes than for R/R homozygotes. The AREase activity in mothers ranged from a low of 19.8 U/ml to a high of 281.4 U/ml (14-fold) and, in newborns, from 4.3–110.7 U/ml (26-fold) (Fig. 2). The range of AREase activity (PON1 levels) from the lowest newborn to the highest mother was 4.3–281.4 U/ml (65-fold). The mean PON1 levels, as measured by AREase, were similar across *PON1*₁₉₂ genotypes for all mothers (Q/Q = 152

Fig. 1



PON1 status plot for mothers and newborns, determined as described in the Methods section. Open circles, data points for PON1_{Q192} homozygous mothers; closed circles, data points for PON1_{Q192} homozygous newborns; closed squares, data points for heterozygous mothers; open squares, data points for heterozygous newborns; open triangles, data points for PON1_{R192} homozygous mothers; closed triangles, data points for PON1_{R192} homozygous newborns.


Fig. 2



Individual data points for arylesterase activities (AREase) in mothers (solid circles) and newborns (open circles) for each PON1₁₉₂ genotype as indicated. Means are indicated by the crossbars.


U/ml, Q/R = 144 U/ml and R/R = 152 U/ml). The mean PON1 levels were also similar across genotypes in newborns (Q/Q = 31 U/ml, Q/R = 36 U/ml and R/R = 43 U/ml), with the newborn PON1_{R192} homozygotes having somewhat higher average AREase activity than the PON1_{Q192} homozygotes and the newborn heterozygotes expressing intermediate levels. However, these AREase activity differences between genotype groups were not statistically significant ($P = 0.13$). Although

Fig. 3



Individual data points for paraoxonase activities in mothers (solid circles) and newborns (open circles) for each PON1₁₉₂ genotype as indicated. Means are indicated by crossbars.

Fig. 4



Plot of arylesterase activities (AREase) of newborn/mother pairs ($r = 0.47$, $P = 0.013$).

newborns had on average four-fold less plasma PON1 than mothers, individual levels in mother-child pairs were modestly correlated ($r = 0.47$, $P = 0.013$) (Fig. 4).

Estimation of the range of sensitivity to organophosphate exposure

Because both PON1₁₉₂ alloforms hydrolyse DZO with the same catalytic efficiency [34], these data predict a range of sensitivity to DZO exposure of 26-fold in newborns and 14-fold in mothers, with a range of 65-fold from the most sensitive newborn to the most resistant mother. An average four-fold difference in sensitivity to DZO exposure is predicted between mothers and newborns.

Estimation of the range of sensitivity to CPO exposure is more complex, due to the different catalytic efficiencies of the two PON₁₉₂ alloforms for CPO hydrolysis. However, reasonable estimates of relative sensitivity can be made by taking advantage of data generated in the mouse model system, where the mouse *PON1* gene was replaced with either human *hPON1*_{Q192} or human *hPON1*_{R192} and colonies were established that expressed these alloforms at the same levels in plasma. Transgenic mice expressing *hPON1*_{Q192} were found to be 2- to 2.5-fold as sensitive to CPO exposure compared to mice expressing *hPON1*_{R192} [48], due to the higher catalytic efficiency of human PON₁₉₂ for hydrolysis of CPO [30]. This two- to 2.5-fold difference in sensitivity, taken together with the 65-fold difference in plasma AREase levels yield an estimated 131–164-fold range in CPO sensitivity between the PON_{1Q/Q192} homozygous newborn with the lowest PON1 level and the PON_{1R/R192} homozygous mother with the highest PON1 level.

Discussion

To our knowledge, this is the first study where PON1 status [39,41,52] was determined for a large cohort of mothers and their newborns using the two-substrate assay. By plotting rates of DZO hydrolysis against rates of PO hydrolysis, the three Q192R phenotypes were clearly separated in the mothers and provided the relative PON1 levels for each individual in this population. However, we found that for 17% of the newborns, with low PON1 levels, PCR analysis was required to assign position 192 genotypes, although the PON1 status analysis did provide their cord blood PON1 levels. Although some of the earlier studies examined only PON1 status with the two substrate diazoxon/paraoxon assay/analysis, this study included a measurement of rates of phenylacetate hydrolysis (AREase), allowing comparison of PON1 levels across *PON1*₁₉₂ genotypes. Another study examining the relationship of PON1 levels, exposure and head circumference in a large cohort of 404 births also made use of AREase activity to determine PON1 levels [55], representing a major improvement over the many studies that examined *PON1* genotypes alone [49]. They noted a relationship between low PON1 levels, exposure and smaller head circumference.

The AREase values provided an estimate of the relative sensitivity to organophosphorus compounds whose in-vivo catalytic efficiencies of hydrolysis are not affected by the *PON1*₁₉₂ polymorphism (e.g. DZO). The range of AREase values in the mothers was very similar to the range observed in another study of Hispanic farm workers in Washington State where a 13-fold variability was observed [28]. The large range of AREase values observed in this study in the newborns (26-fold) predicts a broad variability in sensitivity to organophosphate exposure even among newborns. Surprisingly, some of the new-

borns had higher PON1 levels than some of the adults. Thus, the individuals predicted to be highly sensitive to organophosphate exposure include most of the newborns as well as some of the mothers. The studies reported here confirm earlier observations of lower PON1 activities in neonates compared to adults [23–27,56] and are also consistent with a recent report where neonates had 2.6–4.6-fold lower PON1 levels, as assessed by the AREase assay, compared to mothers, in three ethnic groups in New York City [23].

Among the many activities of the multifunctional HDL-associated enzyme PON1, hydrolysis of DZO and CPO is important in providing protection against exposure to DZ/DZO and CPS/CPO. Organophosphate exposures are appropriately considered as mixed exposures to the parent compounds and their oxon forms because most, if not all, exposures include oxon residues [57,58]. Because the rate of cholinesterase inactivation by CPO is at least three orders of magnitude higher than that of its parent compound (CPS) [59], a very small percentage of oxon form in an exposure is significant.

To date, most of the animal model studies with genetically modified mice have examined CPO/CPS exposures [34,38,46–48,60], although some recent studies were carried out on DZ/DZO exposures in PON1 null mice and PON1 null mice injected with each of the purified human PON₁₉₂ alloforms [34]. The results obtained in these earlier studies have provided important insights and predictions with respect to individual variability in sensitivity to CPO/CPS and DZ/DZO exposures. Li *et al.* [34] examined the relative in-vivo catalytic efficiencies of *hPON1*_{Q192} and *hPON1*_{R192} in PON1 null mice reconstituted with purified human PON₁₉₂ or PON_{1Q192}. They found that under in-vivo physiological conditions, injection of either PON₁₉₂ alloform provided equivalent protection against DZO exposure, whereas injected human PON_{1R192} provided significantly better protection against CPO exposure than did PON_{1Q192}.

A more recent study by Cole *et al.* [48], using transgenic strains of mice that expressed either PON_{1Q192} or PON_{1R192} at nearly equivalent levels, verified the results of Li *et al.* [34] from the enzyme injection studies in the PON1 null mice. The mice expressing PON_{1R192} were approximately two- to 2.5-fold more resistant to CPO exposure than were the *PON1* null mice. Mice expressing PON_{1Q192} were nearly as sensitive as the *PON1* null mice to CPO exposures. The IC₅₀ value (exposure level for 50% inhibition) for CPO inhibition in the *hPON1*_{Q192} transgenic mice was approximately 1.1 mg/kg dermal exposure versus approximately 2.2 mg/kg in the *hPON1*_{R192} mice. The levels of PON1 in the '*hPON1* humanized' transgenic mice are relevant to the studies reported here. The transgenic mice expressed levels of

hPON1 of 30–45 U/ml AREase (corrected for baseline AREase activity in PON1 null mice) [48], which are comparable to the average levels of PON1 in the Latino newborns described in the present study.

Cole *et al.* [24] examined the time course of appearance of PON1 in the plasma of individual children and also in mice expressing the two human PON1₁₉₂ alloforms under control of the human 5' and 3' regulatory sequences. In children, PON1 reached plateau levels at 6–24 months of age whereas expression of human hPON1 in mice under control of the human PON1 5' regulatory region followed the mouse time course of PON1 appearance, peaking at PND21. Moser *et al.* [62] observed that, although AREase activity in rats peaked at 3–4 weeks of age as observed by Li *et al.* [61], resistance to CPS exposure continued to increase up to at least 90 days, corresponding to a continued increase in plasma carboxylesterase activity. Although there is very little carboxylesterase activity in human serum, individual variation in carboxylesterase levels in human liver microsomes has been reported [63].

Levels of PON1 activities reported in other studies provide insights into the range of sensitivities expected. The hydrolytic activities in plasma of the wild-type C57Bl/6J mice used in our earlier studies were approximately 1800 U/l for chlorpyrifos oxonase (CPOase) and approximately 3500 U/l for diazoxonase (DZOase) [60]. These activities are in the low range relative to those reported for human populations. The ranges of reported activities in another adult Hispanic population were 2145–13 540 U/l for CPOase and 2174–23 316 U/l for DZOase [28]. It is worth noting that PON1 null mice are approximately ten-fold more sensitive to DZO [34] and CPO [46] exposure than wild-type mice. Based on what is known about the catalytic efficiencies of organophosphate hydrolysis and the large variability of PON1 levels observed in this and other populations, mothers and newborns homozygous for PON1_{Q192} with very low plasma levels of this alloform are predicted to be a subpopulation uniquely vulnerable to adverse effects from DZ/DZO exposures and especially CPS/CPO exposures. One particular concern would be exposure of a mother with very low PON1 status carrying a fetus that had not yet developed the capacity for self-protection against organophosphate exposure. PON1_{Q192} homozygous mothers with very low PON1 status are also predicted to be unable to pass on to their offspring a PON1 allele that would be protective against exposure. The father of course would contribute one *PON1* allele to the child; however, it would need to be a high expressing allele and would take from 6 months to 2 years following birth to be fully protective.

Although the usual precautions need to be observed in extrapolating data generated in an animal model system

to predict outcomes in human exposures, genetically manipulating a single gene in an inbred strain of mice is nonetheless informative. PON1 appears to be the only major enzyme in the plasma of both mice and humans that hydrolyzes chlorpyrifos oxon and diazoxon, as can be seen from the colinearity of the plots of rates of hydrolysis of one substrate against others in human populations [28] and from the dramatic consequences of deleting this gene in mice [34,46]. Replacing the mouse *PON1* gene with each of the human *PON1*₁₉₂ alleles provides a better means of extrapolating between the two species, even though other pathways may contribute somewhat to the detoxication. Even among humans, there is variability in levels and efficiencies of other contributory detoxication enzymes, such as carboxylesterases and cytochromes P450.

Several factors are likely to contribute to the risk of adverse health effects due to exposures to CPS/CPO and DZ/DZO: (i) the level of exposure; (ii) the percentage of oxon residue or other toxic derivatives in the exposure; (iii) the level of enzyme (measured by AREase assay); (iv) the catalytic efficiency of an individual's PON1 to detoxify organophosphate metabolites (determined by the Q192R polymorphism); and (v) the as yet uncharacterized genetic variability of the cytochromes P450, carboxylesterases and other enzymes that participate significantly in the detoxication of these organophosphorus compounds.

Additional research is needed to determine the relative contributions of PON1, cis- and trans-acting factors that modulate PON1 levels, cytochromes P450 and other enzymes, as well as the genetic and environmental factors that modulate the levels of enzymes involved in detoxication of parent compounds and their respective oxons. A recently reported method of haplotyping using emulsion PCR should be useful in understanding cis factors involved in regulation of protein levels [64]. Research is also needed to determine the exposure levels of specific residues (trichlorophenol, CPS, CPO, diethylphosphate and other toxic metabolites) required for modelling [48,65] the effect of PON1 status on the consequences of exposures.

Evaluation of PON1 status is not only important for determining risk of organophosphate exposure, but also for understanding the role of PON1 in modulating other risks associated with the variability of normal physiological functions of PON1, as well as the role of PON1 in the metabolism of other xenobiotics, including drugs [66]. Multiple results indicating a prominent role for PON1 status in risk for vascular disease have been published previously [40,46,67,68]. More recently, PON1 has been shown to inactivate the quorum sensing signal secreted by *Pseudomonads* [66,69]; however, data demonstrating the

in-vivo importance of this activity have not yet been reported.

In summary, the range of variability of PON1 status observed in this study, taken together with data from 'humanized mice' expressing hPON1_{Q192} or hPON1_{R192} in place of mouse PON1, predict a 65-fold variability in DZO sensitivity and a 131–164-fold range in sensitivity to CPO exposure in this population, with an average four-fold difference in sensitivity to DZO exposure and an average eight-to ten-fold variability in sensitivity to CPO between groups of mothers and their newborns. These data predict that most, if not all, newborns, as well as a subpopulation of adults, will exhibit significantly increased sensitivity to organophosphate exposure. These findings highlight the significance of understanding the susceptibility of young children to organophosphate exposure and developing science-based risk standards for pesticide regulation as required by the 1996 Food Protection Act.

Acknowledgements

We gratefully acknowledge CHAMACOS staff, students and community partners, and especially the CHAMACOS participants and their families, without whom this study would not have been possible. We thank Katie Kogut and Erin Weltzien for their help with statistical analyses and Dr Toby Cole for helpful comments and suggestions.

References

- Barr D, Bravo R, Weerasekera G, Caltabiano L, Whitehead R. Concentrations of dialkyl phosphate metabolites of organophosphorus pesticides in the US population. *Environ Health Perspect* 2004; **112**: 186–200.
- Hill RH Jr, Head SL, Baker S, Gregg M, Shealy DB, Bailey SL, et al. Pesticide residues in urine of adults living in the United States: reference range concentrations. *Environ Res* 1995; **71**:99–108.
- Loewenherz C, Fenske RA, Simcox NJ, Bellamy G, Kalman D. Biological monitoring of organophosphorus pesticide exposure among children of agricultural workers in central Washington State. *Environ Health Perspect* 1997; **105**:344–353.
- Simcox NJ, Camp J, Kalman D, Stebbins A, Bellamy G, Lee IC, Fenske R. Farmworker exposure to organophosphorus pesticide residues during apple thinning in central Washington State. *Am Ind Hyg Assoc J* 1999; **60**: 752–761.
- Berkowitz GS, Obel J, Deych E, Lapinski R, Godbold J, Liu Z, et al. Exposure to indoor pesticides during pregnancy in a multiethnic, urban cohort. *Environ Health Perspect* 2003; **111**:79–84.
- Whyatt RM, Barr DB, Camann DE, Kinney PL, Barr JR, Andrews HF, et al. Contemporary-use pesticides in personal air samples during pregnancy and blood samples at delivery among urban minority mothers and newborns. *Environ Health Perspect* 2003; **111**:749–756.
- Eskenazi B, Bradman A, Gladstone EA, Jaramillo S, Birch K, Holland NT. CHAMACOS: a longitudinal birth cohort study: lessons from the fields. *J Child Health* 2003; **1**:3–27.
- Bradman A, Eskenazi B, Barr DB, Bravo R, Castorina R, Chevrier J, et al. Organophosphate urinary metabolite levels during pregnancy and after delivery in women living in an agricultural community. *Environ Health Perspect* 2005; doi: 10.1289/ehp.7894 [Epub ahead of print].
- Bradman A, Barr DB, Claus Henn BG, Drumheller T, Curry C, Eskenazi B. Measurement of pesticides and other toxicants in amniotic fluid as a potential biomarker of prenatal exposure: a validation study. *Environ Health Perspect* 2003; **111**:1782–1789.
- Castorina R, Bradman A, McKone TE, Barr DB, Harnly ME, Eskenazi B. Cumulative organophosphate pesticide exposure and risk assessment among pregnant women living in an agricultural community: a case study from the chamacos cohort. *Environ Health Perspect* 2003; **111**: 1642–1648.
- Fenske RA, Lu C, Barr D, Needham L. Children's exposure to chlorpyrifos and parathion in an agricultural community in central Washington State. *Environ Health Perspect* 2002; **110**:549–553.
- USEPA 2000. Chlorpyrifos revised risk assessment and agreement with registrants. <http://www.epa.gov/pesticides/op/chlorpyrifos/agreement.pdf>
- USEPA 2001. Diazinon revised risk assessment and agreement with registrants. <http://www.epa.gov/pesticides/op/diazinon/agreement.pdf>
- Sherman JD. Organophosphate pesticides – neurological and respiratory toxicity. *Toxicol Ind Health* 1995; **11**:33–39.
- Eskenazi B, Bradman A, Castorina R. Exposures of children to organophosphate pesticides and their potential adverse health effects. *Environ Health Perspect* 1999; **107** (Suppl 3):409–419.
- Young JG, Eskenazi B, Gladstone EA, Bradman A, Pedersen L, Johnson C, et al. Association between in utero organophosphate pesticide exposure and abnormal reflexes in neonates. *Neurotoxicology* 2005; **26**:199–209.
- Garcia SJ, Seidler FJ, Slotkin TA. Developmental neurotoxicity of chlorpyrifos: targeting glial cells. *Env Toxicol Pharmacol* 2005; **19**: 455–461.
- Moser VC, Padilla S. Age- and gender-related differences in the time-course of behavioral and biochemical effects produced by oral chlorpyrifos in rats. *Toxicol Appl Pharmacol* 1998; **149**:107–119.
- Sheets L. A consideration of age-dependent differences in susceptibility to organophosphorus and pyrethroid insecticides. *Neurotoxicology* 2000; **21**:57–63.
- Padilla S, Buzzard J, Moser VC. Comparison of the role of esterases in the differential age-related sensitivity to chlorpyrifos and methamidophos. *Neurotoxicology* 2000; **21**:49–56.
- Zheng Q, Olivier K, Won YK, Pope CN. Comparative cholinergic neurotoxicity of oral chlorpyrifos exposures in preweanling and adult rats. *Toxicol Sci* 2000; **55**:124–132.
- Pope CN, Chakraborti TK, Chapman ML, Farrar JD, Arthun D. Comparison of *in vivo* cholinesterase inhibition in neonatal and adult rats by three organophosphorothioate insecticides. *Toxicology* 1991; **68**:51–61.
- Chen J, Kumar M, Chan W, Berkowitz G, Wetmur JG. Increased influence of genetic variation on PON1 activity in neonates. *Environ Health Perspect* 2003; **111**:1403–1409.
- Cole T, Jamps RL, Walter BJ, Arndt TL, Richter RJ, Shih DM, et al. Expression of human paraoxonase (PON1) during development. *Pharmacogenetics* 2003; **13**:357–364.
- Ecobichon DJ, Stephens DS. Perinatal development of human blood esterases. *Clin Pharmacol Ther* 1973; **14**:41–47.
- Mueller RF, Hornung S, Furlong CE, Anderson J, Giblett ER, Motulsky AG. Plasma paraoxonase polymorphism: a new enzyme assay, population, family, biochemical, and linkage studies. *Am J Hum Genet* 1983; **35**:393–408.
- Augustinsson K, Barr M. Age variation in plasma arylesterase activity in children. *Clin Chim Acta* 1963; **8**:568–573.
- Davies HG, Richter RJ, Keifer M, Broomfield CA, Sowalla J, Furlong CE. The effect of the human serum paraoxonase polymorphism is reversed with diazoxon, soman and sarin. *Nat Genet* 1996; **14**:334–336.
- Furlong CE, Costa LG, Hassett C, Richter RJ, Adler DA, Distech CM, et al. Human and rabbit paraoxonases: purification, cloning, sequencing, mapping and role of polymorphism in organophosphate detoxification. *Chem Biol Interact* 1993; **87**:35–48.
- Hassett C, Richter RJ, Humbert R, Chapline C, Crabb JW, Omiecinski CJ, Furlong CE. Characterization of cDNA clones encoding rabbit and human serum paraoxonase: the mature protein retains its signal sequence. *Biochemistry* 1991; **30**:10141–10149.
- Sorenson RC, Bisgaier CL, Aviram M, Hsu C, Billecke S, La Du BN. Human serum paraoxonase/arylesterase's retained hydrophobic N-terminal leader sequence associates with HDLs by binding phospholipids: apolipoprotein A-I stabilizes activity. *Arterioscler Thromb Vasc Biol* 1999; **19**:2214–2225.
- Humbert R, Adler DA, Distech CM, Hassett C, Omiecinski CJ, Furlong CE. The molecular basis of the human serum paraoxonase activity polymorphism. *Nat Genet* 1993; **3**:73–76.
- Adkins S, Gan KN, Mody M, La Du BN. Molecular basis for the polymorphic forms of human serum paraoxonase/arylesterase: glutamine or arginine at position 191, for the respective A or B allozymes. *Am J Hum Genet* 1993; **52**:598–608.
- Li WF, Costa LG, Richter RJ, Hagen T, Shih DM, Tward A, et al. Catalytic efficiency determines the in-vivo efficacy of PON1 for detoxifying organophosphorus compounds. *Pharmacogenetics* 2000; **10**:767–779.
- Brophy VH, Jamps RL, Clendenning JB, McKinstry LA, Jarvik GP, Furlong CE. Effects of 5' regulatory region polymorphisms on paraoxonase (PON1) expression. *Am J Hum Genet* 2001; **68**:1428–1436.

- 36 Deakin S, Leviev I, Brulhart-Meynet M-C, James RW. Paraoxonase-1 promoter haplotypes and serum paraoxonase: a predominant role for polymorphic position 107, implicating the Sp1 transcription factor. *Biochem J* 2003; **372**:643–649.
- 37 Suehiro T, Nakamura T, Inoue M, Shiinoki T, Ikeda Y, Kumon Y, *et al.* A polymorphism upstream from the human paraoxonase (PON1) gene and its association with PON1 expression. *Atherosclerosis* 2000; **150**:295–298.
- 38 Li WF, Costa LG, Furlong CE. Serum paraoxonase status: a major factor in determining resistance to organophosphates. *J Toxicol Environ Health* 1993; **40**:337–346.
- 39 Richter RJ, Furlong CE. Determination of paraoxonase (PON1) status requires more than genotyping. *Pharmacogenetics* 1999; **9**:745–753.
- 40 Jarvik GP, Rozek LS, Brophy VH, Hatsukami TS, Richter RJ, Schellenberg GD, Furlong CE. Paraoxonase (PON1) phenotype is a better predictor of vascular disease than is PON1 (192) or PON1 (55) genotype. *Arterioscler Thromb Vasc Biol* 2000; **20**:2441–2447.
- 41 Jarvik GP, Jampsa R, Richter RJ, Carlson CS, Rieder MJ, Nickerson DA, Furlong CE. Novel paraoxonase (PON1) nonsense and missense mutations predicted by functional genomic assay of PON1 status. *Pharmacogenetics* 2003; **13**:291–295.
- 42 Leviev I, Deakin S, James RW. Decreased stability of the M54 isoform of paraoxonase as a contributory factor to variations in human serum paraoxonase concentrations. *J Lipid Res* 2001; **42**:528–535.
- 43 Main AR. The role of A-esterase in the acute toxicity of paraoxon, TEPP and parathion. *Can J Biochem Physiol* 1956; **34**:197–216.
- 44 Costa LG, McDonald BE, Murphy SD, Omenn GS, Richter RJ, Motulsky AG, Furlong CE. Serum paraoxonase and its influence on paraoxon and chlorpyrifos-oxon toxicity in rats. *Toxicol Appl Pharmacol* 1990; **103**: 66–76.
- 45 Furlong CE, Richter RJ, Seidel SL, Costa LG, Motulsky AG. Spectrophotometric assays for the enzymatic hydrolysis of the active metabolites of chlorpyrifos and parathion by plasma paraoxonase/ arylesterase. *Anal Biochem* 1989; **180**:242–247.
- 46 Shih DM, Gu L, Xia YR, Navab M, Li WF, Hama S, *et al.* Mice lacking serum paraoxonase are susceptible to organophosphate toxicity and atherosclerosis. *Nature* 1998; **394**:284–287.
- 47 Li WF, Furlong CE, Costa LG. Paraoxonase protects against chlorpyrifos toxicity in mice. *Toxicol Lett* 1995; **76**:219–226.
- 48 Cole T, Walter B, Shih D, Tward A, Lusi AJ, Timchalk C, *et al.* Toxicity of chlorpyrifos oxon in a transgenic mouse model of the human paraoxonase (PON1) Q192R polymorphism. *Pharmacogenet Genom* 2005; **15**: 589–598.
- 49 Brophy VH, Jarvik GP, Furlong C. PON1 polymorphisms. In: Costa LG, Furlong C, editors. *Paraoxonase (PON1) in health and disease: basic and clinical aspects*. Boston, Massachusetts: Kluwer Academic Press; 2002. pp. 53–77.
- 50 DPR. *Pesticide Use Reporting 2001 Summary Data: Department of Pesticide Regulation*. Sacramento, California: California Environmental Protection Agency; 2002.
- 51 Eskenazi B, Harley K, Bradman A, Weltzien E, Jewell NP, Barr DB, *et al.* Association of in utero organophosphate pesticide exposure and fetal growth and length of gestation in an agricultural population. *Environ Health Perspect* 2004; **112**:1116–1124.
- 52 Richter R, Jampsa R, Jarvik GP, Costa LG, Furlong CE. Determination of paraoxonase 1 (PON1) status and genotypes at specific polymorphic sites. *Curr Protoc Toxicol* 2004; **4**:12.1–4.12.19.
- 53 Blatter-Garin M-C, Abbot C, Messmer S, Mackness MI, Durrington P, Pometta D, James RW. Quantification of human serum paraoxonase by enzyme-linked immunoassay: population differences in protein concentrations. *Biochem J* 1994; **304**:549–554.
- 54 StataCorp. *Stata Statistical Software: release 8.0*. College Station, Texas: Stata Corporation; 2003.
- 55 Berkowitz GS, Wetmur JG, Birman-Deych E, Obel J, Lapinski RH, Godbold JH, *et al.* In-utero exposure, maternal paraoxonase activity and head circumference. *Environ Health Perspect* 2004; **112**:338–391.
- 56 Burlina A, Michielin E, Galzinga L. Characteristics and behaviour of arylesterase in human serum and liver. *Eur J Clin Invest* 1977; **7**:17–20.
- 57 Yuknavage KL, Fenske RA, Kalman DA, Keifer MC, Furlong CE. Simulated dermal contamination with capillary samples and field cholinesterase biomonitoring. *J Toxicol Env Health* 1997; **51**:35–55.
- 58 California EPA 2002. Pesticide use reporting 2001 Summary data. http://www.cdpr.ca.gov/docs/pur/pur01rep/01_pur.htm
- 59 Huff R, Abou-Donia MB. *In vitro* effect of chlorpyrifos oxon on muscarinic receptors and adenylate cyclase. *Neurotoxicology* 1995; **16**:281–290.
- 60 Li W-F. *Dissertation: development of a mouse model for studying paraoxonase*. Seattle, Washington: University of Washington, School of Public Health and Community Medicine, Department of Environmental Health; 1993.
- 61 Li W-F, Matthews C, Distech CM, Costa LG, Furlong CE. Paraoxonase (Pon1) gene in mice: sequencing, chromosomal location and developmental expression. *Pharmacogenetics* 1997; **7**:137–144.
- 62 Moser VC, Chanda SM, Mortensen SR, Padilla S. Age- and gender-related differences in sensitivity to chlorpyrifos in the rat reflect developmental profiles of esterase activities. *Toxicol Sci* 1998; **46**:211–222.
- 63 Hosokawa M, Endo T, Fujisawa M, Hara S, Iwata N, Sato Y, Satoh T. Interindividual variation in carboxylesterase levels in human liver microsomes. *Drug Metabol Dispos* 1995; **23**:1022–1027.
- 64 Wetmur JG, Kumar M, Zhang L, Palomeque C, Wallenstein S, Chen J. Molecular haplotyping by linking emulsion PCE: analysis of paraoxonase 1 haplotypes and phenotypes. *Nucleic Acids Res* 2005; **33**:2615–2619.
- 65 Timchalk C, Poet TS, Hinman MN, Busby AL, Kousba AA. Pharmacokinetic and pharmacodynamic interaction for a binary mixture of chlorpyrifos and diazinon in the rat. *Toxicol Appl Pharmacol* 2005; **205**:31–42.
- 66 Draganov DI, Teiber JF, Speelman A, Osawa Y, Sunahara R, La Du BN. Human paraoxonases (PON1, PON2, and PON3) are lactonases with overlapping and distinct substrate specificities. *J Lipid Res* 2005; **46**:1239–1247.
- 67 Mackness B, Davies GK, Turkie W, Lee E, Roberts DH, Hill E, *et al.* Paraoxonase status in coronary heart disease: are activity and concentration more important than genotype? *Atheroscler Thromb Vasc Biol* 2001; **21**:1451–1457.
- 68 Mackness M, Mackness B. Paraoxonase 1 and atherosclerosis: is the gene or the protein more important? *Free Rad Biol Med* 2004; **37**:1317–1323.
- 69 Ozer EA, Pezzulo A, Shih DM, Chun C, Furlong C, Lusis AJ, *et al.* Human and murine paraoxonase 1 are host modulators of *P. aeruginosa* quorum-sensing. *FEMS Microbiol Lett* 2005; in press.