




Photoelectron Resonance Capture Ionization Aerosol Mass Spectrometry: A New Method for the Study of Organic Aerosols

OVERVIEW

Carbonaceous Particles


- Atmospheric particulate matter (including carbonaceous particles) responsible for poor visibility on humid days:



Comparison of low and high particulate days in Burlington, Vermont


Source: <http://www.hazcam.net/burlington.html>

Contribution to Particulate Composition




Source: <http://www.hazcam.net/burlington.html>

- Significant fraction of total atmospheric particulate matter contains carbon in Eastern U.S.
- Product of **fossil fuel** combustion
 - Complex mixture of compounds
 - Health hazard
 - Undergo active heterogeneous chemistry in atmosphere
- Secondary organic aerosol formation
 - Photochemical oxidative products of (primarily) gaseous emissions from trees generally less volatile than emitted molecules
 - Oxidized products condense to form ultrafine particles



- Atmospheric Processing
 - Particles undergo heterogeneous chemical reactions with atmospheric gas phase species
 - Particles act as sink for **reactive gases**, such as OH and ozone, altering the oxidizing capacity of the atmosphere
 - Physical and chemical properties of particles can be affected – possible **climate change implications**
 - Light scattering properties of particles can be affected
 - Can affect cloud formation
 - Can have severe impact on human health, e.g. nitration of polycyclic aromatic hydrocarbons (PAHs):



Ozonolysis of Oleic Acid Particles

- Has become the **model system** for studying complex behavior of **heterogeneous reactions** involving multi-component organic aerosols.
- Organic fraction of total atmospheric particulate matter is significant and may:
 - Impact **radiative properties** of particles
 - Reactively take up atmospheric **trace gases**
 - Pose severe **human health hazards**
- Relevance of model system to atmospheric processes:
 - Oleic acid is common fatty acid in troposphere (1 ng m⁻³ in remote marine environment)
 - Ozonolysis of oleic acid particles may improve their ability to act as **cloud condensation nuclei** and may disrupt the balance of atmospheric **oxidants**.


METHODS

Experimental Approach

- Methods are needed for the **on-line, real-time** analysis of atmospheric **carbonaceous** particles
- Aerosol Mass Spectrometry** is the method of choice for analysis of atmospheric particles in real-time, however, difficulties are encountered with analysis of organics
- Soft ionization by **photoelectron resonance capture ionization (PERCI)** can minimize molecular fragmentation and simplify molecular identification
- Model heterogeneous reaction system, oleic acid particles and ozone gas, studied to evaluate PERCI
- PERCI allows for the identification of previously undetected particle-phase reaction products

Heterogeneous Chemistry

Polydisperse oleic acid (9-octadecenoic acid) particles are introduced to a flow reactor via a glass tube (0.32 cm I.D.), centered within a 2.54 cm ID flow reactor, which acts as the aerosol injector. Ozone is introduced into the flow reactor upstream of the aerosol injector. The ozone exposure time of the particles is dictated by the position of the aerosol injector at a given flow rate of the particle and gas phases. Reacted particles are introduced into the mass spectrometer through a differentially pumped inlet and focused into a beam using a series of aerodynamic lenses.



Photoelectron Resonance Capture Ionization

PERCI operates on the principle of a tunable, laser-induced photoelectric effect to controllably generate low energy electrons (< 1 eV). Ionization occurs when these electrons resonantly attach, through associative and/or dissociative mechanisms, to nearby gas phase molecules. PERCI provides an exceptionally soft and sensitive ionization, which generates primarily intact molecular ions.

A close-up of the PERCI source, located adjacent to the ion extraction grid of the time-of-flight mass spectrometer. Since the analyte molecules must be in the gas phase for efficient ionization, a heated filament is placed in the particle beam to vaporize the particles.


LaFranchi BW and Petrucci GA, J. Am. Soc. Mass Spectrom. 15:424-430, 2003

LaFranchi BW, Zahradis J, and Petrucci GA, Rapid Comm. Mass Spectrom. in press, 2004.


Acknowledgements: In addition to support from the EPA, the author gratefully acknowledges funding from the American Society for Mass Spectrometry, and Vermont EPSCOR. The author would also like to acknowledge James Zahradis for his contribution to this work.

RESULTS

1. Monitoring the Ozonolysis of Oleic Acid




Facilitated Product Identification




Comparison of PERCI (a) with conventional electron impact (b) mass spectrometry of the ozonolysis products of oleic acid particles. The minimal fragmentation in the PERCI spectrum simplifies identification of the reaction products. The 4 major products shown in the generalized reaction scheme above, are detected as the [M-H]⁺ ions.

Generalized Reaction Scheme




Higher Order Oxygenates Identified



A closer look at the high mass region of the product mass spectrum shows the presence of additional minor products adding to a complex series of reactions and intermediates involved in the ozonolysis of oleic acid.

2. Biodiesel Analysis



Analysis of a complex mixture of organics is possible using PERCI since molecular integrity of the analyte components is maintained. Here a liquid sample of **biodiesel** fuel is analyzed by PERCI. From the mass spectrum, a distribution of the carbon chain lengths of the molecular components in the fuel can be determined. The degree of saturation can also be determined.

CONCLUSIONS

- Four major products of ozonolysis of oleic acid particles observed directly
- Previously undetected higher order oxygenates observed as particle phase products
- Chemical and physical implications in the troposphere:
 - Potential source of OH radical
 - Increased hygroscopicity of oxygenated products enhances the propensity of these particles to act as cloud condensation nuclei
 - Intermediates may react with additional atmospheric gas and particle phase components such as formaldehyde or other environmentally relevant carbonyls
- PERCI can be applied to any type of analysis involving complex mixtures of organics, such as biodiesel fuel