

Final Technical Report

Date of Final Report: August 31, 2006

EPA Grant Number: R827352C002

Center Name: Southern California Particle Center and Supersite (SCPCS)

Center Director: John R. Froines

Title: Pro-inflammatory and the Pro-oxidative Effects of Diesel Exhaust Particulate *in Vivo* and *in Vitro*

Investigators: Andre Nel¹, Ning Li¹, Constantinos Sioutas², Arthur Cho¹, John Froines¹

Institutions: ¹University of California–Los Angeles, ²University of Southern California

EPA Project Officer: Stacey Katz/Gail Robarge

Project Period: June 1, 1999–May 31, 2005 (no-cost extension to May 31, 2006)

Period Covered by the Report: June 1, 1999–May 31, 2006

RFA: Airborne Particulate Matter (PM) Centers (1999)

Research Category: Particulate Matter

Topic A: Studies Emphasizing Investigation of the Biological Mechanisms of Particulate Matter (PM) Effects in Relation to PM Physical and Chemical Characteristics

Objective(s) of the Research Project: Research in this project has increased our understanding of the mechanisms by which PM induce adverse health effects. Progress has been made in understanding the oxidative stress pathways by which diesel exhaust particulate (DEP) and ambient PM mediate injury, and has also helped to elucidate the adjuvant effects of DEP in asthma. We will address research findings and conclusions from our work according to three project aims.

Summary of Findings:

Aim 1: To Elucidate the Role of Reactive Oxygen Species (ROS) and Inflammation in PM-Induced Adverse Health Effects *In Vitro* and *In Vivo*

A potential mechanistic link between PM exposures and inflammation involves the generation of ROS and oxidative stress (Li, et al., 2003; Nel, 2005). A number of our studies, described in more detail below have demonstrated that ambient PM and DEP induce ROS production in target cells such as macrophages and bronchial epithelial cells (Li, et al., 2003a; Nel, 2005; Li, et al., 2002a; Li, et al., 2002b; Hiura, et al., 1999; Li, et al., 2003b). DEP were used in our studies as a convenient model for vehicular ultrafine particles (UF), which are a common component of ambient aerosols in urban areas.

We performed a series of *in vitro* and *in vivo* experiments to explore the link between ROS production, oxidative stress and inflammatory tissue injury (Li, et al., 2003a; Nel, 2005; Mingi, et al., 2003). The findings of the studies form the basis for the development of a hierarchical oxidative stress model (Figure 1). The model posits that at lower levels of oxidative stress (Tier 1), there is an induction of phase II enzymes regulated by a genetic response pathway that involves the transcription factor, Nrf2 (Li, et al., 2003a; Nel, 2005; Li, et al., 2002b; Gilmour, et

al., 2006). Nrf2 drives the antioxidant response element (ARE) in the promoter of phase II response genes, leading to the expression of antioxidant and detoxification enzymes (Li, et al., 2004). Treatment of target cells in vitro with DEP, organic DEP extracts or ambient UF induced the expression of heme oxygenase 1 (HO-1), glutathione-S- transferase (GST), NADPH quinone oxidoreductase (NQO1), catalase, superoxide dismutase (SOD), glutathione peroxidase (GPx) and glucuronosyltransferase (UGT) (Li, et al., 2004; Li, et al., 2000). These phase II enzymes protect against oxidative stress injury (Tiers 2 and 3), and a reduced Tier 1 defense could therefore promote PM susceptibility (Li, et al., 2003a; Nel, 2005). In humans, reduced defenses against oxidative stress can result from phase II enzyme polymorphisms, e.g., the GST M1 null genotype which predisposes to the development of asthma and enhanced sensitization to common environmental allergens during nasal DEP challenge (Li, et al., 2003a). Conversely, induction of a phase II response may be a key factor in adaptation to a polluted environment, and may explain why persistent inflammatory changes in the lung are not observed after repeated exposure to low concentrated ambient particle (CAP) levels. Some phase II enzymes, such as HO-1, exert anti-inflammatory effects based on their ability to interfere with pro-inflammatory response pathways (Li, et al., 2000; Li and Nel, 2006).




Figure 1. Hierarchical Oxidative Stress Model

If Tier 1 protection fails, our model proposes that further increase in oxidative stress generates pro-inflammatory responses (Tier 2) or cytotoxic effects (Tier 3) depending upon the level of insult and response capability of the exposed cells (Figure 1). Tier 2 responses are linked to the activation of intracellular signaling pathways which impact cytokine and chemokine production (Li, et al., 2003a). An example is activation of the MAP kinase cascade (Li, et al., 2003a). This cascade is responsible for the expression and activation of AP-1 transcription factors (e.g., c-Jun and C-Fos), which in turn are responsible for the expression of a variety of pro-inflammatory genes, including those encoding for cytokines, chemokines and adhesion molecules. Tier 3 responses in our model involve mitochondrial perturbation by pro-oxidative chemicals (Hiura, et al., 1999; Li, et al., 2003b; Hiura et al., 2000; Xia, et al., 2004). Although the in vivo significance of the mitochondrial pathway is uncertain, we have demonstrated in tissue culture cells that PM-induced interference in one electron transfers in the mitochondrial inner membrane and perturbation of the mitochondrial permeability transition pore (PTP) can contribute to

superoxide generation and the induction of cellular apoptosis (Hiura, et al., 1999; Li, et al., 2003b; Hiura, et al., 2000; Xia, et al., 2004). These effects can be mimicked by organic extracts made from DEP as well as redox cycling quinones and functionalized aromatic hydrocarbons present in the DEP particles (Li, et al., 2002a; Li, et al., 2000). Each tier of oxidative stress is sensitive to the effects of NAC (Li, et al., 2002b; Hiura, et al., 1999; Whitekus, et al., 2002; Xia, et al., 2004).

The principles of a hierarchical oxidative stress response were tested in macrophages and epithelial cell cultures exposed to ambient UF, DEP extracts, or fractionated DEP extracts (Li, et al., 2002a). At the lowest tier of oxidative stress (Tier 1), the expression of catalase, SOD, and HO-1 indicates the involvement of Nrf2-regulated enzymes that can suppress inflammation through their antioxidant activities (Li, et al., 2003a; Li, et al., 2004). This finding was extended by showing that particulate pollutants increase the accumulation of Nrf2 in the nucleus and activate the ARE (Li, et al., 2004). Interestingly, the buildup of Nrf2 in the nucleus is dependent on a prolongation of protein half-life by interference in proteosomal degradation (Li, et al., 2004). Activation of the ERK, p38 and Jun kinase cascades was confirmed by phosphor-proteome analysis (Wang, et al., 2005). To further substantiate the findings, related experiments are now being conducted in vivo, using BAL fluid and lung tissue from PM-exposed animals to find in vivo biomarkers of oxidative stress. These markers could be useful to identify the subsets of the human population susceptible to PM exposure.

While there is still considerable debate about which particle components are responsible for the pro-oxidative and pro-inflammatory effects associated with PM, our work adds to accumulating evidence that transition metals, such as copper, vanadium, chromium, nickel, cobalt and iron, as well as aromatic and polar organic substances play a role in ROS production (Li, et al., 2003a; Li, et al., 2000). The particle backbone could play an important role in acting as a template for single electron transfers reactions, including electron transfer to molecular dioxygen (Figure 2). This could involve redox cycling reactions, as demonstrated by the ability of ambient PM samples to generate superoxide in the presence of dithiothreitol (DTT) (Li, et al., 2003b). DTT oxidation can be assessed by a colorimetric reaction to assay for the content of redox cycling chemicals in urban PM samples (Li, et al., 2003b). In addition, biologically catalyzed oxidation-reduction reactions in the cellular interior, as well as interference in one electron transfers in the mitochondrial inner membrane, contribute to ROS generation (Xia, et al., 2004). In addition to the ability of ROS to damage cellular proteins, DNA and cell membranes, electrophilic PM chemicals such as the quinones can modify cellular proteins by Michael acceptor reactions (Li, et al., 2004). It is likely that this type of reaction leads to Nrf2 release to the nucleus by the covalent modification of its cytosolic chaperone, Keap-1 (Li, et al., 2004). The covalent modification of intracellular and tissue proteins was also confirmed by studying their tyrosylation and carbonylation after in vivo exposure to diesel exhaust particulate (Whitekus, et al., 2002).




Figure 2. Particle-Induced ROS Production by Key Chemical Components

In experiments to characterize the redox cycling chemicals present in PM, silica gel chromatography was used to fractionate organic DEP extracts (Li, et al., 2002a; Li, et al., 2003b; Li, et al., 2000; Xia, et al., 2004). Aliphatic, aromatic and polar chemical fractions were eluted by increasingly polar solvents and tested for reactivity in the DTT assay. The quinone-enriched polar material was more active than the polycyclic aromatic hydrocarbon (PAH)-enriched aromatic fraction. Glutathione depletion in epithelial cells and macrophages is associated with the exposure to DEP extracts and with activity in the DTT assay (Li, et al., 2002a; Li, et al., 2003b; Li, et al., 2000; Xia, et al., 2004). The aliphatic fraction was inactive in these assays.

The relationship between the organic chemical composition and the redox cycling potential of PM that had been noted for diesel particles was confirmed in a study in which UF were compared to coarse particles (C) and fine particles (F) collected in the Los Angeles Basin (LAB) (Li, et al., 2003b). UF were more active than C and F in the DTT assay, and were also more prone to generate oxidative stress in macrophages and epithelial cells (Li, et al., 2003b). Both the *in vitro* and cellular responses showed an excellent correlation with the PAH content of UF (Nel, 2005; Li, et al., 2003b). Another important observation in this study was the ability of UF to lodge in and disrupt the mitochondrial architecture (Li, et al., 2003b). This finding is related to cellular apoptosis and apo-necrosis by a pathway that requires opening of the mitochondrial PTP (Hiura, et al., 2000; Xia, et al., 2004). Functional effects on the PTP and inability to sustain one electron transductions in the mitochondrial inner membrane was confirmed in isolated mitochondrial preparations through the use of calcium-dependent swelling, calcium retention capacity and dissipation of the mitochondrial membrane potential (Xia, et al., 2004). Moreover, UF particle effects could be reproduced by polar and aromatic chemicals fractionated from DEP, while commercial polystyrene nanoparticles were inactive (Xia, et al., 2004). These data demonstrate differential particle toxicity associated with particle size, composition, and subcellular localization.

Aim 2: To Develop a Murine Model for Asthma to Explain the Adjuvant Effects of DEP on Ovalbumin (OVA)-Induced Allergic Inflammation and Airway Hyperreactivity (AHR)

The asthma studies were premised on findings that DEP enhance allergen-specific IgE and TH2 cytokine production in humans and animals (Li, et al., 2003a; Nel, 2005). We demonstrated that aerosolized DEP can enhance OVA-specific IgE production in a murine inhalation model (Whitekus, et al., 2002). The adjuvant effect of DEP could be suppressed by NAC administration (Whitekus, et al., 2002). While adequate for upregulating IgE production, an

important limitation of this model was the inability of DEP to enhance AHR. This likely reflects the fact that oxidizing chemicals lead to efficient IgE gene rearrangement in deposition hotspots (Li, et al., 2003a), but exposure did not exceed the threshold of airway inflammation that is required for AHR (Li, et al., 2003a). DEP-induced AHR has now been achieved through modification of the classical mouse OVA sensitization model, in which sensitization is achieved by intraperitoneal administration of OVA. We developed two new protocols for OVA sensitization (Mingi, et al., 2003). In the low grade sensitization protocol, BALB/c mice received intraperitoneal OVA without alum, followed by challenge with aerosolized OVA $\pm$ DEP two weeks later. In the post-challenge model, DEP was delivered to classically sensitized animals a few days after the OVA challenge. Under both conditions, DEP enhanced airway inflammation to the point of exceeding the AHR threshold (Mingi, et al., 2003). Since these data suggest that low grade airway inflammation is required to elicit AHR, nebulized DEP was administered to mice which have been genetically engineered to overexpress IL-5. These animals exhibit constitutive airway inflammation, and responded to DEP inhalation with increased airway inflammation and AHR (Mingi, et al., 2003).

To demonstrate the role of oxidative stress in the adjuvant effects of DEP in allergic inflammation, we performed an animal study in which mice were sensitized to OVA and co-challenged with aerosolized DEP daily for ten days (Whitekus, et al., 2002). The thiol antioxidants, N-acetylcysteine (NAC) and bucillamine (BUC) were administered intraperitoneally during sensitization. NAC and BUC effectively inhibited the adjuvant effects of DEP in the induction of OVA-specific IgE and IgG1 production (Whitekus, et al., 2002). Furthermore, NAC and BUC prevented the generation of lipid peroxidation and protein carbonylation, an oxidative modification of protein structure, in the lungs of OVA- plus DEP-exposed animals. These findings indicate that NAC and BUC are capable of preventing the adjuvant effects of inhaled DEP and suggest that oxidative stress is a key mechanistic component in the adjuvant effect of DEP. Antioxidant treatment strategies may therefore serve to alleviate allergic inflammation and may provide a rational basis for treating the contribution of particulate matter to asthmatic disease.

Aim 3: Dosimetry and Distribution of Particles

Health effects of exposure to PM are likely to be proportional to the PM dose to critical cells and organs. Tissue dose is influenced by the proportion of inhaled particles that are retained in the lung with each breath. The proposed hierarchical oxidative stress response that occurs in PM target cells has been discussed above. A frequently asked question is how the experimental in vitro DEP concentrations that span the three tiers of oxidative stress (e.g., 1–100 $\mu\text{g}/\text{ml}$ in macrophages) can be understood in terms of tissue concentrations that are achieved in the lung during real-life exposures. One method to reconcile doses used in vitro with in vivo exposures, is to convert ambient PM levels, measured in $\mu\text{g}/\text{m}^3$, to a dose that is deposited on a planar surface and then to compare that to the calculated dose of the DEP that are deposited on a planar tissue culture surface (Li, et al., 2003a). Our calculations resulted in a planar concentration of 0.2–20 $\mu\text{g}/\text{cm}^2$ on the tissue culture dish. This concentration was compared to a theoretical in vivo deposition dose that would occur in the nasopharyngeal (NPR), tracheobronchial (TBR), and alveolar (AVR) regions of the respiratory tract of an adult person exposed to $\text{PM}_{2.5}$ in Rubidoux, California (Li, et al., 2003a). After correction for parameters such as airway anatomy,

nasal breathing, high rates of deposition at bifurcation points, and uneven airflow due to airway obstruction in asthma or chronic obstructive pulmonary disease (COPD), the calculated deposition values for the NPR, TBR and AVR were 204, 2.3 and 0.05 $\mu\text{g}/\text{cm}^2$, respectively, for the Rubidoux scenario. The calculations showed that it is possible to achieve doses in the nose and TBR from ambient exposures that are responsible for the in vitro induction of antioxidant, pro-inflammatory and cytotoxic responses.

References:

Li N, Hao M, Phalen RF, Hinds WC, Nel AE. Particulate air pollutants and asthma: a paradigm for the role of oxidative stress in PM-induced adverse health effects. *Clinical Immunology* 2003a;3:250-265.

Nel A. Air pollution-related illness: biomolecular effects of particles. *Science* 2005;308: 804.

Li N, Wang M, Oberley TD, Sempf JM, Nel AE. Comparison of the pro-oxidative and proinflammatory effects of organic diesel exhaust particle chemicals in bronchial epithelial cells and macrophages. *Journal of Immunology* 2002a;169:4531-4541.

Li N, Kim S, Wang M, Froines J, Siouts C, Nel A. Use of a stratified oxidative stress model to study the biological effects of ambient concentrated and diesel exhaust particulate matter. *Inhalation Toxicology* 2002b;14:459-486.

Hiura TS, Kaszubowski MP, Li N, Nel AE. Chemicals in diesel exhaust particles generate reactive oxygen radicals and induce apoptosis in macrophages *Journal of Immunology* 1999;163:5582-5591.

Li N, Sioutas C, Cho A, Schmitz D, Misra C, Sempf J, Oberley T, Froines J, Nel A. Particulate air pollutants, oxidative stress and mitochondrial damage. *Environmental Health Perspectives* 2003b;111:455-460.

Whitekus M, Li N, Zhang MJ, Wang M, Horwitz M, Nelson M, Brechun SK, Diaz-Sanchez N, Nel D, Thiol AE. Antioxidants inhibit the adjuvant effects of aerosolized diesel exhaust particles in a murine model for ovalbumin sensitization. *Journal of Immunology* 2002;168:2560-2567.

Minqi Hao, Comier S, Wang M, Lee J, Nel A. Diesel exhaust particles exert acute effects on airway inflammation and function in murine allergen provocation models. *The Journal of Allergy and Clinical Immunology* 2003;112:905-914.

Gilmour IM, Jaakkola MS, London SJ, Nel AE, Rogers CA. How the indoor and outdoor environments influence the incidence and severity of asthma. *Environmental Health Perspectives* 2006;114:627-633.

Li N, Alam J, Venkatesan MI, Eiguren-Fernandez A, Schmitz D, Di Stefano EM, Slaughter N, Killeen E, Wang X, Huang A, Wang M, Miguel AH, Cho A, Sioutas C, Nel AE. Nrf2 is a key transcription factor that regulates antioxidant defense in macrophages and epithelial cells:

protecting against the pro-inflammatory and oxidizing effects of diesel exhaust chemicals. *Journal of Immunology* 2004;173:3467-3481.

Li N, Venkatesan MI, Miguel A, Kaplan R, Gujuluva C, Alam J, Nel A. Induction of heme oxygenase-1 expression in macrophages by diesel exhaust particle chemicals and quinones via the antioxidant-responsive element. *Journal of Immunology* 2000;165:3393-3401.

Li N, Nel AE. Role of the Nrf2-mediated signaling pathway as a negative regulator of inflammation: implications for the impact of particulate pollutants on asthma. *Antioxidants & Redox Signaling* 2006;8:88-98.

Hiura TS, Li N, Kaplan R, Horwitz M, Seagrave JC, Nel AE. The role of a mitochondrial pathway in the induction of apoptosis by chemicals extracted from diesel exhaust particles. *Journal of Immunology* 2000;165:2703-2711.

Wang M, Xiao GG, Li N, Xie Y, Loo JA, Nel AE. Use of a fluorescent phosphoprotein dye to characterize oxidative stress-induced signaling pathway components in macrophage and epithelial cultures exposed to diesel exhaust particle chemicals. *Electrophoresis* 2005;26(11):2092-2108.

Xia T, Korge P, Weiss JN, Li, Venkatesen I, Sioutas C, Nel A. Quinones and aromatic chemical compounds in particulate matter (PM) induce mitochondrial dysfunction: implications for PM-induced oxidative stress and toxicity. *Environmental Health Perspectives* 2004;112:1347-1358.

Supplemental Keywords: NA

Relevant Web Sites: <http://www.scpcs.ucla.edu>